

Exercise metabolism and adaptation in skeletal muscle

Jonathon A. B. Smith¹, Kevin A. Murach², Kenneth A. Dyar^{3,4} & Juleen R. Zierath^{1,5,6} ✉

Abstract

Viewing metabolism through the lens of exercise biology has proven an accessible and practical strategy to gain new insights into local and systemic metabolic regulation. Recent methodological developments have advanced understanding of the central role of skeletal muscle in many exercise-associated health benefits and have uncovered the molecular underpinnings driving adaptive responses to training regimens. In this Review, we provide a contemporary view of the metabolic flexibility and functional plasticity of skeletal muscle in response to exercise. First, we provide background on the macrostructure and ultrastructure of skeletal muscle fibres, highlighting the current understanding of sarcomeric networks and mitochondrial subpopulations. Next, we discuss acute exercise skeletal muscle metabolism and the signalling, transcriptional and epigenetic regulation of adaptations to exercise training. We address knowledge gaps throughout and propose future directions for the field. This Review contextualizes recent research of skeletal muscle exercise metabolism, framing further advances and translation into practice.

Sections

Introduction

Skeletal muscle fibre types and subcellular characteristics

Acute exercise metabolism in skeletal muscle

Skeletal muscle responses to acute exercise

Skeletal muscle adaptations to long-term exercise

Future directions

¹Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden. ²Molecular Mass Regulation Laboratory, Exercise Science Research Center, Department of Health, Human Performance and Recreation, University of Arkansas, Fayetteville, AR, USA. ³Metabolic Physiology, Institute for Diabetes and Cancer, Helmholtz Diabetes Center, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany. ⁴German Center for Diabetes Research (DZD), Neuherberg, Germany. ⁵Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden. ⁶Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. ✉e-mail: juleen.zierath@ki.se

Introduction

As a primary site of nutrient storage, energy use and locomotion, skeletal muscle is central to the impact of physical activity on human health. Periods of inactivity reduce skeletal muscle insulin sensitivity and oxidative capacity¹, contributing towards impaired systemic metabolic flexibility² and an increased risk for cardiometabolic disease³. Physical activity offers a degree of protection against the deleterious effects of sedentary behaviour on whole-body metabolism⁴: larger volumes⁴ and more vigorous types⁵ of physical activity, such as formal exercise, probably convey additional benefit.

Bouts of exercise rapidly sensitize skeletal muscle to hormones (Supplementary Table 1) and nutrients. A single (or ‘acute’) exercise session directly increases skeletal muscle transport of amino acids⁶ and glucose⁷. These effects appear somewhat specific to the contracted musculature, and they enhance postprandial muscle protein synthesis⁸ and insulin-stimulated glucose disposal^{9,10} in the recovery period after exercise. Consistent exercise training (over weeks, months and years) further augments skeletal muscle mass^{11–13}, peripheral insulin-sensitivity¹², maximal oxygen consumption ($\dot{V}O_{2\max}$)^{12–15} and strength^{12,13}. $\dot{V}O_{2\max}$ (ref. 16) and strength¹⁷ are well-known predictors of mortality, and training for their improvement through endurance and resistance exercise (Box 1) reduces mortality risk in a manner that is most effective when both training modalities are performed within the same exercise programme¹⁸.

Although the association between regular physical activity and health span has been realized since antiquity, recent methodological advances have allowed the field of exercise physiology to progress towards comprehensive systems-level profiling of the complex molecular interplay that occurs with exercise^{19,20}. These advancements have enabled better mechanistic understanding of the cause-and-effect relationships underlying exercise adaptation and associated health benefits.

In this Review, we provide a contemporary summary of the role of skeletal muscle in response to exercise. First, we address the major cellular makeup of skeletal muscle, highlighting fibre type properties, and the importance of sarcomeric and mitochondrial networks. Next, we discuss skeletal muscle metabolism during acute exercise and the influence of select modifiers such as intensity and timing on this biology. Finally, we address underlying mechanisms of exercise-induced skeletal muscle adaptation and consider differences between training modalities that may facilitate distinct and complementary responses beneficial for human performance and health.

Skeletal muscle fibre types and subcellular characteristics

The human body contains >600 skeletal muscles mainly comprising long contractile cells called muscle fibres. The complex architecture of these fibres (Fig. 1a) provides clues into the intricacies of muscle function and homeostatic control. In this section we focus on the contractile and metabolic properties of different muscle fibre types and examine how these attributes are supported by interacting networks of sarcomeres and mitochondria.

Contractile, metabolic and myonuclear properties of muscle fibre types

Human muscles of the torso and limbs express three main fibre types, with slow-to-fast contractile properties in the following order: slow-oxidative myosin heavy chain (MyHC) type I (encoded by *MYH7*), fast oxidative-glycolytic (intermediate) MyHC type IIA (encoded by *MYH2*)

and fast-glycolytic MyHC type IIX (formerly known as type IID; encoded by *MYH1*)²¹. Muscle fibres have classically been ‘typed’ according to metabolic – oxidative versus glycolytic enzyme – profiles^{22,23} and the predominant abundance^{22,24} or ATPase activity²³ of MyHC isoforms. MyHC are the motor proteins of myofibril thick filaments and determine important aspects of muscle function, such as maximum shortening velocity²⁵. Yet, the full extent of fibre type characteristics (Supplementary Fig. 1) depends upon matching the excitation–contraction coupling machinery and ATP provision to MyHC activity^{14,21}. This coordinated expression is a product of α -motor unit innervation and the transcriptional synchrony of resident myonuclei^{26,27}.

During differentiation (myogenesis) (Supplementary Box 1), a combination of intracellular forces ‘squeeze’ centrally located myonuclei to the fibre periphery²⁸, where they reside in a generally ordered pattern of ‘domains’ in mature muscle. Discrete populations of myonuclei serve specialized roles within fibres, including those governing the neuromuscular junction, myotendinous junction^{26,29,30} (the interface between muscle and tendon) and proprioceptive muscle spindles³⁰. Likewise, the function of ‘body’²⁶ or ‘canonical’²⁹ myonuclei (constituting $\geq 90\%$ of total myonuclei)²⁶ is to direct fibre type specificity in muscle, and they can be identified in mouse muscle by their *Myh* isoform signature^{26,29,30}. *Myh*-positive myonuclei from type I versus type II fibres possess unique chromatin accessibility and transcription factor motif enrichment profiles that are distinct from one another and from myonuclei in other cellular compartments, including the myotendinous junction²⁶. This could underlie specific transcription factor-driven myonuclear programmes^{26,30} that are partially responsible for regionalized gene expression in muscle cells^{26,30,31} and for the co-expression of specific calcium (Ca^{2+})-handling, sarcomeric and metabolic apparatus^{14,21,26,31} so that fibre type contractile (also known as ‘twitch’) and metabolic properties generally align (Supplementary Fig. 1b).

The myonuclei in fibres from endurance-trained younger and older individuals are more spherical, contain greater lamin A (LMNA) deposition and are stiffer and less deformable than myonuclei in untrained counterparts³². These structural and mechanical modifications may facilitate the transduction of cytoskeletal forces towards the nucleus and improve myonuclear resilience against contractile damage³². As such, exercise training-induced myonuclear remodelling could have important implications for muscle adaptation and integrity across the lifespan.

In general, human muscles often express a greater proportion of slower-twitch fibres than those of other species^{22,24}, and human fibres are slower-contracting than orthologous fibre types in most mammals, including rats and mice²⁴. Similarly, type IIA fibres are the most oxidative fibre type in rodent muscle, whereas type I fibres are most oxidative in humans²² (Supplementary Fig. 1). Such differences might contribute to the lack of conformity between the transcriptomes of human and mouse muscle after acute or chronic resistance exercise³³ and should be considered when inferring human relevance from animal physiology. Moreover, the properties of muscle can vary markedly among individuals^{23,34}, biological sexes²³ (Supplementary Box 2), anatomical locations³⁵ and during ageing^{34,36} (Box 2). For example, discrete spatial metabolomic differences within fibre types have been observed in mouse muscle³⁷, and age-associated mitochondrial impairments may induce a glycolytic shift in human fibres without a corresponding change in MyHC³⁶. Collectively, this cautions against the use of MyHC as a strict marker of metabolism and vice versa. Future inquiry should better define how covariates – including biological sex, social gender, biological versus chronological age, metabolic health and

Box 1

Different exercise (sub)types and training response heterogeneity in humans

Exercise (sub)types

Exercise can be broadly classified as resistance, cardiorespiratory, balance and flexibility-based. Flexibility and balance are important aspects of physical fitness contributing to fall prevention in older individuals²⁸². However, discussion herein is dedicated to iterations of resistance and cardiorespiratory exercise, which are the main focus of this Review.

Resistance exercise

Traditional resistance exercise entails repetitions of dynamic concentric (muscle-shortening) and eccentric (muscle-lengthening) contractions against external load and is an effective intervention to increase skeletal muscle mass and strength^{249,283}. The total amount (volume), frequency and intensity of exercise are inherently linked training variables that impact adaptation and performance. For resistance exercise, volume is often reported as the number of times (or 'sets') a particular muscle group is trained per week and is the main stimulus for muscle accrual²⁸⁴. Twelve to 20 weekly sets is sufficient to maximize hypertrophy²⁸⁵, and the frequency of exercise can be adjusted to disperse training volumes according to personal preference²⁸⁶.

Intensity of resistance exercise is generally normalized to a percentage of the maximum load that an individual can lift (expressed as the percentage of 1 repetition maximum (RM))²⁸⁷ or how close an exercise set is taken to momentary muscular failure (in other words, the inability to complete the concentric portion of a movement)²⁸⁸. Equal gains in muscle mass can be made irrespective of load-intensity (<30% 1 RM to ≥80% 1 RM)²⁸⁷ when sets are taken proximal to failure²⁸⁸. Thus, increasing the number of repetitions against a fixed load or increasing the load lifted for a fixed number of repetitions are both viable progression strategies to promote muscle hypertrophy²⁸⁹. Conversely, absolute strength (1 RM)²⁸⁷, tendon stiffness²⁹⁰ and running economy (that is, the metabolic cost at a given velocity of submaximal running)²⁵⁴ are improved more by high-load resistance training. Collectively, this implies that implementing various loading strategies could represent the best approach for attaining the breadth of muscle-related adaptations to resistance training.

The stress imposed by a single bout of resistance exercise upregulates amino acid transporters and sensors in muscle at least in part through activating transcription factor 4 (ATF4)⁶. Synergizing with the mammalian target of rapamycin (mTOR) (see the section 'Skeletal muscle responses to acute exercise') (Fig. 3), this sensitizes muscle to amino acids for ≥24–48 h after resistance exercise^{6,8}, and dietary protein intakes of ≥1.2–1.6 g per kg body mass daily modestly complement the hypertrophic response to resistance training²⁹¹.

Endurance and high-intensity cardiorespiratory exercise

Cardiorespiratory exercise has numerous subtypes that are associated with training-induced improvements in maximal oxygen

consumption ($\dot{V}O_{2max}$)^{292–294}. Approaches used to standardize cardiorespiratory exercise are plentiful (reviewed in ref. 295) and most often include the allocation of intensity against fixed percentages of maximal power (\dot{W}_{max}), velocity (\dot{V}_{max}), heart rate (HR_{max}) or $\dot{V}O_{2max}$. However, the validity of normalizing cardiorespiratory exercise to maximal parameters has been questioned^{295,296}, as this can produce dissimilar physiological and metabolic perturbations between individuals²⁹⁶. Moving forward, studies should strive to establish methods of standardization that provoke homeostatic disturbances consistent with distinct exercise intensity domains.

Endurance (also known as aerobic) exercise is typically considered to be a continuous bout of formal activity performed at low (<50%), moderate (~50–79%) or high intensities (≥80%) of $\dot{V}O_{2max}$ (ref. 292). High-intensity cardiorespiratory exercise can be further divided into high-intensity interval exercise (HIE) and sprint interval exercise (SIE), both of which constitute several bursts of higher-intensity effort interspersed with periods of low-intensity active recovery. High-intensity intervals are conducted at ≥80% of $\dot{V}O_{2max}$, whereas sprint intervals are supramaximal or 'all-out'²⁹². Comparison of cardiorespiratory exercise suggests that training within the high-intensity range is more effective^{292,293} or equally as effective²⁹⁴ at increasing $\dot{V}O_{2max}$ and requires less training volume than moderate intensities to do so²⁹⁴. HIE and SIE training also improve endothelial function to greater extents, whereas moderate intensities favour long-term glycaemic control (lower glycated haemoglobin A_{1c} (HbA_{1c}))²⁹². Further intricacies are seen between interval training subtypes. Eight weeks of HIE training improved cardiac stroke volume, $\dot{V}O_{2max}$ and endurance (3 km) performance²⁹⁷. By contrast, an equivalent period of SIE training potentiated anaerobic capacity and sprinting (300 m) performance²⁹⁷.

Training variables might also discretely regulate muscle mitochondria (see the section 'Skeletal muscle adaptations to long-term exercise'). Hence, combinations of endurance exercise, HIE and SIE are recommended to maximize both health and performance benefits. Consistent with this, just 1–3 h per week of moderate or high-intensity cardiorespiratory exercise could lower mortality risk, and incorporating resistance exercise confers added protection¹⁸.

Interindividual variation in exercise adaptation

The magnitude of adaptation to resistance²⁹⁸ and endurance^{299,300} training differs substantially between individuals. In part, this probably stems from the aforementioned challenges regarding exercise standardization^{295,296}, but also from genetic³⁰⁰ and environmental interactions that converge to produce the heterogeneous molecular responses that are observed systemically¹⁹ and in muscle¹⁷⁷ after a common exercise bout.

Interindividual difference in exercise trainability has led to the concept of 'responders' and 'non-responders'. Increasing the volume of fixed-intensity exercise^{301,302} or the intensity of

(continued from previous page)

fixed-volume exercise³⁰² can somewhat attenuate exercise non-response, with larger volumes of higher-intensity exercise perhaps doing so most effectively³⁰². However, lower responders still require a greater training stimulus and time commitment to achieve results comparable to those of more-responsive trainees^{301,302}.

training status – interact to determine the full spectrum of muscle characteristics.

Hybrid fibre types

Most canonical myonuclei within the same fibre display coordinated transcription of a single *Myh* isoform in mouse muscle^{26,27}. However, a minority of fibres are hybrid^{26,27} (reviewed in ref. 38), containing myonuclei that express two or more *Myh* pre-mRNA genes in the same nucleus^{26,29} and/or in different nuclei across the fibre length²⁶. Thus, the regional distribution of MyHC can vary between muscle biopsy sites³⁹ and along a single fibre^{26,40}.

In human vastus lateralis muscle, <10%^{22,41} to 40%³⁸ of fibres can be hybrid types. ‘True’ non-transitioning or non-regenerating hybrid fibres have metabolic enzyme²² and single-fibre contractile (force-velocity producing)^{25,42} properties between those of their co-expressed MyHC isoforms, providing further functional nuance to the slow-oxidative to fast-glycolytic continuum in the following order of slowest and most oxidative to fastest and most glycolytic: type I → I/IIA → IIA → IIA/IIX → IIX^{22,25,42}. In adult mice, hybrid fibres are most common in the slow-twitch soleus^{26,27}, and the abundance of hybrid fibres is not altered by denervation in this muscle group²⁶. By contrast, sciatic nerve transection or deletion of *Six1* – a gene encoding the transcription factor homeobox protein SIX1 driving the fast-glycolytic phenotype in muscle⁴³ – increases hybrid fibre content in the typically fast-twitch extensor digitorum longus^{26,27}. Hence, anatomical position²⁷, innervation^{26,27} and transcription factor profiles²⁶ might coalesce to coordinate *Myh* expression in muscle.

For a given muscle group, a greater proportion of hybrid and pure type IIX fibres seems indicative of sedentary behaviour⁴¹. Alternatively, exercise tends to reduce hybrid fibre content^{41,44,45}, promoting a shift away from type IIX fibres towards slower myosin types^{13,39,41}. Consequently, pure type IIX fibres are rare in humans^{22,24,39,41} and account for <1% of the vastus lateralis fibre pool in healthy individuals⁴⁰. As discussed in Supplementary Box 1, *Myh* expression in mouse muscle is regulated by competitive promoter–enhancer interactions^{27,46}. Exercise epigenetically modifies chromatin accessibility (see the section ‘Skeletal muscle responses to acute exercise’), and resistance exercise increases MyHC-specific protein synthesis⁴⁷. Yet, it is still unclear how established regulators of fibre type switching, such as peroxisome proliferator-activated receptor- γ coactivator 1 α (PGC1 α , also known as PPARGC1 α)⁴⁸ and nuclear factor of activated T cells, cytoplasmic 1 (NFATc1)⁴⁹, combine with *Myh*-promoter–enhancer dynamics to confer physical activity-dependent fibre type transitions. Further single-myonuclei RNA and chromatin profiling, alongside isolated fibre spatial transcriptomic and proteomic approaches, should extend the understanding of how gene expression and cellular phenotype are regulated among the many nuclei of a syncytial muscle fibre.

As the field of sports genetics continues to grow and be refined, it should provide actionable knowledge for the efficient personalization of training programmes. In the meantime, it is important to emphasize that low responders for a given parameter, such as $\dot{V}O_{2\max}$, often improve in other outcome measures²⁹⁹. Therefore, exercise remains beneficial for all who are able to partake.

The myofibrillar matrix

The sarcomere is the basic contractile unit of skeletal muscle, and in-series assembly of sarcomeres forms myofibrils. Although myofibrils were initially viewed as single, tube-like structures organized in parallel within fibres, later evidence suggested that myofibrils instead form branching networks that could act in tandem with the cytoskeleton to facilitate lateral-force transmission⁵⁰. Using focused ion beam-scanning electron microscopy, recent studies have built upon this hypothesis showing that myofibrils indeed form a nonlinear lattice of sarcomeres^{51–53} connected across the width and length of the muscle cell through three branching subtypes⁵¹ (Supplementary Fig. 2a). In mice, the frequency of sarcomere branching decreases during early-to-late postnatal development, but increases again in adult slow-oxidative soleus fibres⁵¹. In comparison to soleus muscle, fast-glycolytic gastrocnemius fibres exhibit a different branching morphology (preferring myofilament transfer over sarcomere splitting) and utilize fewer total sarcomeric connections⁵¹. Inducing sarcomere connections, such as through gene manipulation in *Drosophila*, reduces the myofibril cross-sectional area⁵². Thus, myofibrillar matrix assembly appears specific to functional demand, and its organization in fast-twitch muscle may support the greater size²² and power²⁴ of these fibres.

The relevance of a nonlinear network of sarcomeres is implied by its structure – providing an elegant mechanism for both longitudinal and lateral force transmission from muscles to bones through tendons^{50,51}. A linked configuration of myofibrils could also minimize the impact of localized sarcomere damage in muscle and thereby increase the robustness of the contractile machinery across an entire fibre. Further interrogation of the function and regulation of the myofibrillar matrix in health, disease and exercise adaptation will be enlightening.

Mitochondrial complexity in muscle

Sustained muscle contraction requires a continuous supply of ATP, which is mostly derived from mitochondrial oxidative phosphorylation (OXPHOS) during submaximal⁵⁴ or longer-duration high-intensity exercise⁵⁵ (see below; Fig. 2). Depending on fibre type, around 2–10%⁵⁶ of muscle volume is filled by distinct subpopulations of subsarcolemmal (also known as peripheral) and intermyofibrillar mitochondria, differing in structure, function and localization^{56–59} (Fig. 1).

Subsarcolemmal mitochondria often cluster together in the sarcoplasmic space between myofibrils and the sarcolemma^{56–59}, and they dedicate more of their volume to cristae (folds of the inner membrane) and matrix^{57,58}. These globular mitochondria extend deep into the myofibrillar space and physically join with adjacent intermyofibrillar mitochondria through electron-dense intermitochondrial junctions^{57,58}, forming the mitochondrial reticulum^{56–59} (Fig. 1b,c). Compared to subsarcolemmal mitochondria, intermyofibrillar mitochondria are more complex in morphology⁵⁹ and physically interact

Glossary

Adrenoceptor

Adrenergic transmembrane G-protein-coupled receptors (GPCRs) that mediate the actions of the endogenous catecholamines adrenaline and noradrenaline. There are nine subtypes of adrenoceptors: α_{1A} , α_{1B} , α_{1D} , α_{2A} , α_{2B} , α_{2C} , β_1 , β_2 and β_3 . The α_{2A} and α_{2C} adrenoceptors regulate presynaptic neurotransmitter release from central adrenergic and peripheral sympathetic nerves.

Ergogenic

A performance-enhancing effect.

Hypertrophy

Typically refers to an increase in the cross-sectional area (or radial growth) of muscle fibres, resulting in gains of skeletal muscle mass in response to mechanical loading activities, such as resistance exercise.

Ketone body

A lipid-derived, water-soluble, organic compound produced in the liver that can be used as an alternative energy source by extra-hepatic tissues — predominantly the brain, but also heart and skeletal muscle.

Maximal oxygen consumption

($\dot{V}O_{2max}$). The maximum volume of oxygen ($\text{ml kg}^{-1} \text{min}^{-1}$) that can be inspired and utilized during exhaustive exercise, such that the value ($\dot{V}O_2$) plateaus despite increasing workloads. $\dot{V}O_{2max}$ is a measure of aerobic or cardiorespiratory fitness and is commonly used to standardize exercise intensity for clinical trials (for example, % of $\dot{V}O_{2max}$).

Mitophagy

A specific form of lysosome-dependent catabolism (autophagy), through which damaged mitochondria are selectively removed. Mitophagy of the mitochondrial reticulum has an essential role in maintaining cellular energy homeostasis.

Muscle spindles

Structures embedded in most mammalian skeletal muscles that continuously relay proprioceptive information regarding muscle length and movement to the central nervous system. Muscle spindles consist of intrafusal muscle fibres enclosed within a capsule layer and are distinct from the extrafusal muscle fibres discussed in this Review.

Non-esterified fatty acids

(NEFAs). A metabolic substrate utilized by muscle at rest and in an intensity-dependent manner during exercise.

Peak-twitch torque

The force produced by muscle (through a moment arm) evoked by a single electrical stimulation from, for example, applied electrodes.

Phosphagen system

A rapid energy-producing pathway comprising the ATP regenerating adenylate kinase ($\text{ADP} + \text{ADP} \rightleftharpoons \text{ATP} + \text{AMP}$) and creatine kinase ($\text{CrP} + \text{ADP} \rightleftharpoons \text{ATP} + \text{Cr}$) reactions. Of these reactions, creatine kinase has a greater capacity for ATP resynthesis in muscle due to the availability of creatine phosphate stores.

Proprioceptive

Able to sense intrinsic information regarding bodily position and locomotion. The primary proprioceptive sensory organ of the body is the muscle spindle.

Proton-motive force

The proton electrochemical gradient in mitochondria consisting of an electrical charge gradient (also known as the 'membrane potential') and a pH gradient. The proton-motive force is generated by the proton-pumping action of respiratory complexes across the inner mitochondrial membrane and couples substrate oxidation to ATP generation.

Transverse tubules

(T-tubules). Invaginations in the sarcolemmal membrane that insert between myofibrils. T-tubules tightly associate with two terminal cisternae (calcium-releasing regions) of the sarcoplasmic reticulum, forming the 'triads', which are essential for excitation-contraction coupling.

Voluntary force production

The conscious or 'voluntary' production of muscle force (in other words, not triggered by exogenous electrical stimulation).

with the myofibrillar matrix^{53,56,57}, sarcoplasmic reticulum (SR) and intermyofibrillar lipid droplets^{53,56,58} (Fig. 1b).

In humans^{53,59,60} and mice^{53,56}, intermyofibrillar mitochondrial (sub)networks are uniform, but their organization varies between muscle types^{56,60}. In fast-glycolytic extensor digitorum longus fibres of mice, intermyofibrillar mitochondria wrap around the I-bands of sarcomeres, perpendicular to the contraction axis, whereas slow-oxidative soleus fibres contain larger subnetworks of connected intermyofibrillar mitochondria, arranged in grids that surround myosin like a cage⁵⁶. The positioning of mitochondria in the intermyofibrillar space also directly impacts the structure of adjacent sarcomeres⁵³. A greater proportion of mitochondria at the Z-disc reduces the cross-sectional area of sarcomeres in this region by bending (or 'curving') peripheral myosin filaments, causing heterogenous myosin-myosin spacing along the sarcomere length⁵³. However, any detrimental impact on fibre contractility might be offset by the relative preservation of cross-sectional area and myofilament spacing towards the sarcomere centre (M-line)⁵³ (the structure of an individual sarcomere is shown in Supplementary Fig. 2d).

Subsarcolemmal mitochondria are proximal to nutrient-delivering capillaries⁵⁶⁻⁵⁹ and have a greater abundance of proton-motive force driving electron transport chain (ETC) complex IV⁵⁷. As such, this subpopulation is thought to specialize in membrane potential

generation for subsequent transfer through the reticulum, into the intermyofibrillar mitochondrial network⁵⁷. Here, intermyofibrillar mitochondria, possessing higher ATP synthase (ETC complex V) expression⁵⁷ and higher surface area-to-volume ratios⁵⁹, could utilize this potential energy to support rapid ATP production and diffusion to myofibrillar ATPases⁵⁶. Hence, much like an electrical power grid, the mitochondrial reticulum may serve to efficiently disperse energy across the fibre⁵⁶⁻⁵⁸.

Within mitochondria, ETC complexes form higher-order structures termed 'supercomplexes'. The maximal oxygen consumption of permeabilized fibres was positively correlated with the preferential redistribution of complexes III and IV into supercomplexes in the muscle of elderly individuals after 4 months of endurance training⁶¹. Alternatively, 6 weeks of high-intensity exercise improved muscle mitochondrial respiration and ETC enzymatic activity in the absence of supercomplex alterations in young adults¹⁵. Supercomplexes are clearly important structural features of the ETC, but whether they support exercise adaptation beyond their stoichiometric relationship with mitochondrial abundance is yet to be determined. Furthermore, the assembly and stability of supercomplexes depends on the integrity of mitochondrial cristae^{62,63}, which can be regulated by SR stress-induced signalling based on eukaryotic translation initiation factor 2 α (EIF2 α) kinase 3 (PERK, also known as EIF1AK3), EIF2 α




Fig. 1 | Skeletal muscle fibre ultrastructure. **a**, Location of mitochondrial subpopulations and energy stores in muscle fibres. Skeletal muscle is composed of layers of connective tissue and fascicles (also known as muscle bundles). Fascicles contain organized arrangements of individual syncytial muscle fibres, each covered by an endomysium, or basal lamina, which is anchored to the fibre membrane (also known as the sarcolemma). Muscle stem cells, termed ‘satellite cells’, reside within this sarcolemma-basal lamina ‘niche’ (Supplementary Fig. 3). Specialized components, such as sodium/potassium pumps (Na^+/K^+ -ATPase), triads (consisting of transverse tubule and sarcoplasmic reticulum (SR)) and proteins of the myofibrils (long arrangements of connected sarcomeres) enable fibre contraction through the process of excitation–contraction coupling and sliding filament theory (Supplementary Fig. 2). Free ATP in muscle is limited^{69,70}, and fibres possess additional energy depots to maintain contractile activity, including creatine phosphate, glycogen and intramyocellular lipids (Box 3). Glycogen granules are nonuniformly distributed between intramyofibrillar, intermyofibrillar and subsarcolemmal pools^{35,267,268}. Alternatively, intramyocellular lipids are stored in lipid droplets (LDs) found predominantly at central (intermyofibrillar) but also peripheral (subsarcolemmal) regions within healthy muscle fibres^{76,78}. During submaximal⁵⁴ and longer-duration high-intensity interval⁵⁵ exercise most ATP in muscle is regenerated by mitochondrial oxidative phosphorylation (OXPHOS) (see the section ‘Acute exercise metabolism in skeletal muscle’) (Fig. 2). **b**, Spatial distribution of the mitochondrial reticulum within muscle fibres. Human muscle comprises

three main fibre types^{14,21,22,24,36}, type I (marked by *MYH7* expression), type IIA (with *MYH2* expression) and type IIX (expressing *MYH1*). Differences in mitochondrial protein content^{14,21} and mitochondrial network configuration^{56,60} between fibre types directly impacts muscle metabolism and function. Muscle mitochondria form an interconnected reticulum^{56–60} that enables swift and efficient distribution of potential energy from subsarcolemmal (also known as peripheral) mitochondria to intermyofibrillar mitochondria (IMF), deep within the fibre^{56–58}. The positioning of mitochondria in the intermyofibrillar space influences the structure of adjacent sarcomeres, resulting in variable cross-sectional areas and myofilament spacing at different regions across the sarcomere length⁵³. The branching morphology of IMF also accommodates functional interactions with nearby cellular components, such as the sarcoplasmic reticulum and intermyofibrillar lipid droplets^{53,56,58}. In oxidative mouse muscle, ~20% of all IMF are connected to lipid droplets, which may facilitate efficient ATP production and distribution⁵⁶. **c**, Adjacent mitochondria form networks and share energy potential through the intermitochondrial junction (IMJ). Analogous to circuit breakers, intermitochondrial junctions split the reticulum into smaller subnetworks, permitting swift separation of defective mitochondria before their removal through mitophagy⁵⁸. In this way, intermitochondrial junctions provide a dynamic layer of quality control, rapidly rewiring the mitochondrial reticulum through healthy network components to sustain muscle function⁵⁸. C, cytochrome c; CoQ, coenzyme Q; IMM, inner mitochondrial membrane; IMS, intermembrane space; OMM, outer mitochondrial membrane.

and activating transcription factor 4 (ATF4) (PERK–EIF2 α –ATF4)⁶³. Endurance-trained athletes have greater cristae density in mitochondria of type I fibres⁶⁴, and PERK and ATF4 proteins are increased in muscle ~48 h after acute resistance exercise⁶⁵. However, cristae structure was unchanged after 10 weeks of moderate-intensity endurance training in sedentary individuals with obesity⁶⁴. Thus, further study is required to distinguish the impact of exercise on mitochondrial cristae remodelling.

Moving forward, better understanding of mitochondrial networks, subpopulations and ETC configurations could benefit strategies to combat age-associated decline in muscle function (Box 2) and mitochondria-related diseases. Focus should be directed towards the heterogeneous mitochondrial populations in human muscle, which are smaller than those in mice⁵⁹.

Acute exercise metabolism in skeletal muscle

Relative to weight, the basal thermogenesis of muscle is lower than that of most other organs⁶⁶ because myosin is maintained in disordered relaxed and super-relaxed states characterized by slow and extremely slow ATP kinetics, respectively⁶⁷ (Supplementary Figs. 2b,c). Upon contraction, mechanosensing rapidly initiates myosin conformational change from relaxed to active⁶⁸, and muscle ATP consumption increases dramatically during short-duration exhaustive exercise^{69,70}. As free ATP in muscle (~20–25 mmol per kg dry mass) is only sufficient to sustain maximal exercise for <2 s (refs. 69,70), continued contractile activity requires ATP resynthesis from a combination of intramuscular energy stores (Fig. 1a and Box 3) and circulating substrates, such as glucose and non-esterified fatty acids (NEFAs)^{71,72} (Fig. 2).

In this section we detail the metabolic responses that enable muscle to match the considerable demands of acute exercise. We also address how this substrate–energy pairing facilitates the integral role of muscle in exercise-mediated inter-organ communication (Box 4) and touch upon the interaction between exercise metabolism and biological rhythms.

Muscle substrate utilization is exercise intensity-dependent

Muscle fibres are part of a functional ‘motor unit’ comprising an α -motor neuron and the muscle fibres innervated by its axon. The force generated by a muscle depends on both the number of activated motor units – and thus fibres – and the rate at which motor units discharge action potentials once recruited (known as rate coding) (reviewed in ref. 73). Stimulation of motor units conforms to the size-orderly principle of recruitment, such that smaller units are activated first, followed sequentially by larger units as contraction intensifies⁷³. Consequently, the higher power outputs achieved during progressively demanding physical activity relies on the stimulation of a greater number of motor units, a larger proportion of the muscle fibre pool, and therefore the recruitment of more type II fibres.

ATP consumption per unit of time is ~2.5–4-fold higher in type II fibres than in type I fibres⁷⁴, and the maximal rate of ATP resynthesis is fastest through oxygen-independent (anaerobic) versus oxygen-dependent pathways, in the following order of descending speed: anaerobic phosphagen system (including adenylate kinase and creatine kinase (CKM)) and glycolysis reactions provide the fastest supply of ATP, followed by carbohydrate oxidation and finally NEFA oxidation. Human type II fibres are hence enriched with creatine phosphate (CrP) and glycogen energy depots⁷⁵ and contain higher levels of adenylate kinase²¹, glycolysis and glycolysis metabolic machinery^{14,21}. Conversely, type I fibres are more abundant in peroxisomes¹⁴, mitochondria^{14,21,76,77} and intramyocellular lipids (IMCLs)^{76–78}, consistent with their slower ATP turnover⁷⁴ (Supplementary Fig. 1b).

Together, this indicates that the contribution of specific energy systems and substrates to working muscle is mainly a function of the neuromuscular activation required to match the intensity of the exercise being performed. This understanding is envisaged within the ‘crossover concept’ (reviewed in ref. 79), which describes the larger relative contribution of NEFA oxidation towards whole-body energy expenditure at low-to-moderate exercise intensities, with an incremental and necessary ‘switch’ towards preferential (oxygen-dependent and

Box 2

Effects of ageing on skeletal muscle and the benefits of exercise

The loss of total muscle mass³⁰³, fibre number³⁴ and type II fibre size^{34,36,250} becomes most evident at ≥ 50 years of age^{34,303} and is inherently linked to perturbations in muscle metabolism. Mitochondrial content is lower in type I and type II fibres from older individuals than from younger individuals³⁶, and diminished muscle oxidative capacity is linked to mobility decline among adults ≥ 60 years of age^{84,304}. Glycolytic enzymes and chaperone proteins of myofibrillar quality control are also decreased in type II versus type I fibres from individuals > 65 years of age³⁶. Together with an age-dependent decline of *MYH1* and *MYH2* mRNA expression (encoding myosin heavy chain IIX isoform (MyHC-IIX) and MyHC-IIA, respectively)⁴⁷, this disconnect between muscle metabolic and contractile apparatus could contribute towards the reduced type II fibre size^{34,36,250} that predominantly underlies detriments in muscle force and power with advanced age³⁰⁵ (reviewed in ref. 306). Older humans and aged mice can further accrue subsets of senescent muscle fibres³⁰⁷ and resident mononuclear cells (for example, satellite³⁰⁸, myeloid^{307–309} and fibroadipogenic progenitor^{307,308} cells). The number of senescent cells is typically low in resting muscle but increases after resistance exercise³¹⁰ and injury^{308,309} irrespective of age. The defective clearance and potential accumulation of senescent cells during ageing^{308,309} might impair muscle regeneration^{308,309}, hypertrophy (especially of fast-twitch glycolytic fibres)³¹⁰, strength^{307,311} and maximal mitochondrial respiration³¹¹, in part through a *Cdkn1a*-driven transcriptional programme^{307,311}.

Exercise can mitigate several aspects of muscle ageing and improves systemic insulin sensitivity¹². Lifelong endurance exercisers have a greater density of muscle mitochondria^{312,313} and a more complex and connected mitochondrial reticulum³¹³ relative to less-active elderly counterparts. This is coincident with higher protein levels of inner mitochondrial membrane fusion-factor optic atrophy 1 (OPA1)³¹³. Additionally, resistance training was shown to offset age-related CpG-site methylations in the mitochondrial

genome³¹⁴, and long-term mixed-modality endurance-type exercise might attenuate methylation events in the promoter regions of important cytoskeletal, sarcomeric, glycolysis, glycogen synthesis and tricarboxylic acid cycle-related genes³¹². These epigenetic events could combine to preserve muscle contractile and metabolic integrity. Furthermore, exercise training-induced changes in plasma apelin (APLN) positively correlated with chair stand, balance and walking (Short Physical Performance Battery) test scores in elderly women and men³¹⁵. This suggests that factors produced from exercising muscle help to mediate the protective effects of physical activity against age-related functional decline (Box 4). APLN released from muscle during exercise can act in an autocrine manner to augment satellite cell-mediated repair, mitochondrial abundance and muscle oxidative capacity³¹⁵, and/or through paracrine mechanisms to stimulate endothelial cell expansion³¹⁶ that could enhance hypertrophy of type II fibres^{248,250} (Supplementary Fig. 3).

Endurance⁶¹, high-intensity interval and concurrent¹² training all increase maximal oxygen consumption ($\dot{V}O_{2max}$) and muscle mitochondrial content in older individuals. Reciprocally, resistance training improves strength^{12,240,317} and fat-free mass¹² but to a lesser extent than seen in young adults²⁴⁰. The attenuated muscle anabolic response with age could be due to blunted ribosomal biogenesis²⁴⁰, reduced activity of amino acid sensors (such as leucyl tRNA synthetase (LARS))³¹⁸ and lower systemic production and local tissue sensitivity to hormones such as testosterone and insulin-like growth factor 1 (IGF1)²⁴⁰ (Supplementary Table 1). Progressive resistance training programmes can further benefit hip and femur bone mineral density in people ≥ 65 years of age³¹⁷, and incorporating power-type resistance exercise (involving explosive concentric movements) may be superior for physical performance outcomes, including 'get-up-and-go' and chair stand tests³¹⁹. Collectively, these findings emphasize the importance of maintaining a diverse physical activity profile across the lifespan.

independent) carbohydrate utilization during exercise at higher levels of mechanical effort^{71,72}.

Metabolic inertia at the onset of exercise

Although the lower metabolic cost of rest⁷² and light exercise⁷¹ is mainly fuelled by the oxidation of circulating NEFAs, further upregulation of OXPHOS ATP provision is delayed at the onset of moderate-to-high intensity physical activity, and a larger oxygen-independent contribution fuels muscle in the first ≤ 30 – 60 s of exercise^{54,80}. This is despite sufficient blood flow^{54,80} and adequate intramuscular oxygen levels for maximal mitochondrial respiration to occur (≥ 0.5 – 2 mmHg) (reviewed in ref. 81).

The lag in oxygen-dependent metabolism at the beginning of exercise may stem from a combination of linked temporal factors, including activation of the mitochondrial matrix enzymes pyruvate dehydrogenase (PDH), 2-oxoglutarate dehydrogenase (OGDC)⁸² and carnitine acetyltransferase (CRAT)^{83,84}; the availability of tricarboxylic acid (TCA)

cycle substrates (pyruvate and glutamate)⁸² and precursors (acetyl groups)^{83–85}, and sarcoplasmic buffering of the interrelated allosteric metabolites creatine and ADP⁸⁶ (Fig. 2a). Indeed, pharmaceutical activation of PDH increased muscle acetyl-carnitine availability and reduced the reliance on phosphagen and glycolytic energy pathways during acute ischaemic exercise⁸⁵. The ability of previous sprinting intervals to prime OXPHOS in subsequent high-intensity work bouts could also occur through better coupling of oxidative substrate delivery to mitochondrial enzymatic activity, although carryover of residual fatigue probably plays an additional role in this scenario by impeding power output^{69,70}.

Oxygen-independent exercise metabolism

During short-duration maximal efforts, accelerated muscle ATP demand is mostly met anaerobically, through rapid stimulation of the phosphagen and glycolytic energy systems^{69,70}. The simplicity and proximity of the sarcoplasmic CKM reaction (Box 3) allows CrP to supply equimolar amounts of high-energy phosphate to ATPases within milliseconds⁸⁷,

which is essential for sprint performance^{69,70}. CrP hydrolysis peaks at the onset of contraction but declines within less than 6 s, and CrP stores can be >90% depleted after around 30 s of intense exercise⁷⁰ compared to only ~20% after 10 min of cycling at ~50% of $\dot{V}O_{2\max}$ (ref. 88).

The initial increase in glycogen breakdown is due to posttranslational modification of glycogen phosphorylase (PYGM) from a less-active *b* form to the constitutively active *a* form by phosphorylase kinase (PHK)⁷⁰. Transients of Ca^{2+} and AMP are probable regulators of PHK and thus the switch to PYGM*a* (reviewed in ref. 89). AMP can simultaneously increase PYGM*b* activity to support maximal rates of glycogenolysis⁷⁰, and the systemic rise of adrenaline during sprint intervals⁶⁹ or endurance exercise^{71,90,91} may help to stabilize the PYGM*a* conformation⁹². Downstream of glycogen and glucose, allosteric regulation and replenishment of nicotinamide adenine dinucleotide (NAD⁺) can stimulate the rate-limiting enzymes phosphofructokinase (PFK) and glyceraldehyde-3-phosphate dehydrogenase (GAPDH), respectively, to promote continued flow through glycolysis (Fig. 2c).

Average power output drops substantially over the final half of a 30-s all-out physical effort, corresponding to the pattern of PYGM activity (reversion from *a* to *b*) and reduced substrate-level phosphorylation⁷⁰. This muscular fatigue might be driven by alterations to the intracellular metabolic milieu. Above a critical threshold of ~15 mmol l⁻¹ (ref. 93), inorganic phosphate can enter the SR and precipitate with Ca^{2+} , thereby impairing Ca^{2+} release and muscle contraction⁹⁴. Additionally, owing to the high activity and near-equilibrium state of lactate dehydrogenase (LDH), the end product of glycolysis is always lactate⁹⁵. After ~3.5–4 min of exhaustive cycling, muscle lactate reaches levels more than twofold higher than resting concentrations⁵⁵, more than fourfold higher following acute resistance exercise⁹⁶, and ≥8-fold⁹⁷ to 30-fold⁶⁹ higher after 10 rounds of short-duration sprinting. The accumulation of lactate (a strong anion) encourages dissociation of water to $HO^- + H^+$, and muscle pH can reach ≤6.5 during strenuous exercise bouts⁵⁵. Lower intramuscular pH could diminish PYGM activity⁷⁰, and H^+ may act on group III (mainly mechanosensitive) and IV (mainly metabosensitive) muscle afferents in the interstitial space⁹³. Once activated, these sensory neurons can feed back, potentially through inhibitory γ -aminobutyric acid type B (GABA_B) receptors⁹⁸, to reduce motor cortex excitability and suppress motoneuronal output⁹³. The severity of peripheral fatigue correlated with the extent of quadriceps activation during self-paced time-trial cycling⁹⁹. However, mechanisms of exercise-induced peripheral and central fatigue are complex, and the major contributory factors are probably specific to modality, intensity and duration (reviewed in ref. 94).

Oxygen-dependent exercise metabolism

Reactive oxygen species production during exercise. In contrast to maximal sprinting for ≤30 s (refs. 69,70), high-intensity cycling lasting ≥3 min derives >70% of energy from OXPHOS, with a post-exercise muscle metabolome enriched for pathways using pyruvate, long-chain NEFAs and amino acids such as alanine, arginine and glutamate⁵⁵. This upregulation of OXPHOS reduces mitochondrial hydrogen peroxide (H₂O₂) emission^{100,101} from ETC complexes I–III¹⁰⁰ and thus mitochondria contribute minimally to reactive oxygen species (ROS) generation in contracting muscle¹⁰². Rather, nicotinamide adenine dinucleotide phosphate (NADPH) oxidases (NOX), located in the sarcolemma and in transverse tubules (T-tubules)¹⁰² and the sarcoplasmic reticulum¹⁰³, appear to be the predominant sources of ROS during contraction^{101,102}. Specifically, NOX2 (ref. 101) and NOX4 (ref. 104) are indispensable for exercise-stimulated sarcoplasmic ROS production in mice.


ROS signalling is implicated in adaptations to endurance and resistance training (reviewed in ref. 105). High-intensity intervals augment NOX4 mRNA in human vastus lateralis ~3–4 h after exercise, and a retrograde NOX4–ROS-axis led to activation of transcriptional regulators – such as nuclear factor erythroid 2-related factor 2 (NRF2, also known as NFE2L2) and PGC1 α – required for mitochondrial biogenesis and the endogenous antioxidant defence response to exercise training in mouse muscle¹⁰⁴ (Fig. 3). Accordingly, blunting oxidative stress through *Nox4* deletion in mice¹⁰⁴ or high-dose vitamin C and E intake in humans¹⁰⁶ attenuates the insulin-sensitizing effects of endurance training. Still, any potential detriment of antioxidant (such as vitamin C and/or vitamin E) supplementation to gains in $\dot{V}O_{2\max}$, lean body mass or human endurance and strength performance seems relatively minor¹⁰⁷.

Glucose uptake and carbohydrate oxidation in exercising muscle.

When workload^{71,72} and (to a lesser extent) duration⁷¹ of moderate-intensity exercise increases, so does the contribution of blood glucose towards total carbohydrate utilization, although energy provision from intramuscular glycogen is always higher^{71,72}. Muscle glucose uptake reaches ~15-fold resting levels at the cessation of non-fatiguing exercise (~50% of $\dot{V}O_{2\max}$) and ~50-fold resting levels at the cessation of exhaustive endurance exercise (~100% of $\dot{V}O_{2\max}$)⁸⁸. This increased rate of blood glucose extraction is supported by greater hepatic glucose output¹⁰⁸, a ≥5–10-fold rise in blood flow⁸⁸, and enhanced perfusion of muscle capillaries¹⁰⁹. As exercise intensifies, these circulatory and microvascular responses collectively maintain relatively constant plasma-to-interstitial glucose concentrations¹¹⁰ and enlarge the area available for nutrient exchange to occur¹⁰⁹. Such events further serve to promote the release of discrete biologically active molecules from muscle²⁰. As discussed in Box 4, many of these muscle-derived ‘exerkines’ or exercise-induced ‘myokines’ are implicated in aspects of local and systemic exercise adaptation (reviewed in ref. 111).

The exocytosis and fusion of glucose transporter 4 (GLUT4)-containing vesicles with the sarcolemmal membrane and T-tubules is essential for contraction-stimulated glucose transport into muscle¹¹². During exercise, translocation of GLUT4 is regulated by a combination of Ca^{2+} , metabolic stimuli and mechanosensitive stimuli that converge on calcium/calmodulin-dependent protein kinase II (CaMKII), 5'-AMP-activated protein kinase (AMPK) and RAS-related C3 botulinum toxin substrate 1 (RAC1) (reviewed in ref. 113), with redundancy between pathways^{113,114}. The role of RAC1 in contraction-induced glucose uptake might be mediated in part by downstream NOX2 activation¹⁰¹ but seems independent of AMPK α_2 (ref. 114) during submaximal treadmill running in mice. Indeed, the catalytic α subunit of AMPK appears dispensable for exercise-stimulated glucose transport in vivo^{115,116}. Instead AMPK α plays a more notable role in post-exercise substrate metabolism¹¹⁷ and insulin sensitivity¹¹⁶ through the upregulation of pyruvate dehydrogenase kinase 4 (PDK4)¹¹⁷, the promotion of RAS-related protein RAB8A-perilipin 5 (PLIN5) lipid droplet-mitochondrial tethering¹¹⁸ and the phosphorylation of TBC1 domain family member 1 (TBC1D1)¹¹⁵ and TBC1D4 (also known as AS160)¹¹⁶.

For glucose to be oxidized in muscle, lactate – simultaneously produced from glycolysis and taken up through sarcolemmal monocarboxylate transporters (MCT1 and MCT4)⁸¹ – must first be converted to pyruvate by sarcoplasmic or mitochondrial LDH (reviewed in ref. 119) (Fig. 2c). Glucose oxidation is higher in men than in women (Supplementary Box 2) and rises with exercise intensity, dietary carbohydrate and muscle glycogen levels, and peri-exercise



carbohydrate intake¹²⁰. Intra-workout consumption¹²¹ or ‘mouth-rinsing’¹²² of exogenous carbohydrates can also benefit exercise performance. Whereas the ingestion of carbohydrates feasibly provides metabolizable substrate¹²⁰ and spares liver glycogen stores during bouts of longer-duration exercise¹⁰⁸, the potential ergogenic effect of mouth-rinsing is most probably achieved through oral receptor afferents that signal centrally to increase voluntary force production¹²³.

Muscle lipid metabolism during exercise. The post-exercise plasma metabolome is distinct between exercise modalities. An acute bout of endurance or resistance exercise differentially regulates discrete subclusters of metabolites across a 3-h post-exercise period, despite similar trends for metabolome recovery over time¹²⁴. Of note, amino acid, nucleotide and carbohydrate (for example, lactate and pyruvate) signatures are prominent after resistance exercise, compared to the lipid-derivative enrichment (for example, various acyl-carnitines and the ketone body β -hydroxybutyrate) after endurance exercise¹²⁴. This illustrates unique physiological challenges posed by specific interventions.

At intensities eliciting peak fat oxidation (~60–65% of $\dot{V}O_{2max}$), the contribution of plasma NEFAs and IMCLs is ~1:1 and roughly equal to total carbohydrate utilization^{71,72}. In mouse muscle, liberation of NEFAs

from lipid droplets is almost entirely dependent¹²⁵ on the redundant enzymes^{125,126} adipose triacylglyceride lipase (ATGL) and hormone-sensitive lipase (HSL). The recruitment of ATGL to PLIN5 at lipid droplets could be enhanced by AMPK-regulated assembly of the RAB8A–PLIN5 tethering complex¹¹⁸. Alternatively, exercise upregulates HSL in an intensity-dependent manner¹²⁷ through additive contraction (Ca^{2+} –protein kinase C (PKC))¹²⁸ and adrenaline (cAMP–PKA)⁹¹-mediated pathways that promote its translocation to lipid droplets¹²⁹. These signalling events augment rates of muscle lipolysis, and IMCL contents of type I and type II fibres can be ~45% and ~20% depleted, respectively, after ~60–180 min of cycling at ~50–75% of $\dot{V}O_{2max}$ (ref. 130). Muscle HSL activity is transient during acute moderate-intensity exercise⁹¹ and its downregulation – through negative feedback from AMPK⁹¹ or allosteric metabolites (for example, long-chain fatty acyl-CoA) – mirrors the greater relative contribution of circulating NEFAs towards whole-body energy expenditure⁷¹.

Circulating NEFA availability is increased during longer bouts of physical activity at low-to-moderate intensity⁷¹ due to greater release from adipose tissue and the improved affinity of triacylglycerol-rich lipoproteins for hydrolysis by lipoprotein lipases (LPL)¹³¹ anchored to the endothelial surface of interstitial capillaries. The utilization of plasma NEFAs is enhanced by contraction-induced sarcolemmal

Fig. 2 | Skeletal muscle metabolism during higher-intensity exercise.

a, Exercise-onset metabolic inertia (red area). Acetyl-carnitine (aCarn) abundance^{83–85} and the acetyl coenzyme A (aCoA)-producing capacities of carnitine acetyltransferase (CRAT)^{83,84} and pyruvate dehydrogenase (PDH)⁸⁵ appear rate-limiting for oxidative adenosine triphosphate (ATP) regeneration at the onset of moderate-high intensity exercise. Contraction-induced calcium (Ca^{2+}) transients promote mitochondrial Ca^{2+} uptake into the matrix space²⁶⁹ through the inner-membrane mitochondrial calcium uniporter (MCU) complex^{270,271}. Increased matrix Ca^{2+} can upregulate PDH²⁷² through activation of its phosphatase (PDP)²⁷³, and can upregulate isocitrate dehydrogenase (IDH)²⁷⁴ and 2-oxoglutarate dehydrogenase (OGDC)^{272,274,275} directly to fine-tune oxidative metabolism via stimulation of the tricarboxylic acid (TCA) cycle⁸². Ca^{2+} kinetics probably precede an allosteric⁸⁶ rise in the $[\text{ADP}]/[\text{P}_i]$ to $[\text{ATP}]$ and $[\text{creatine}]/[\text{P}_i]$ to $[\text{creatine phosphate}]$ ratios in part because ADP and creatine (Cr) are buffered by the adenylate kinase (not shown) and creatine phosphate (CrP) shuttle reactions. Thus, synchrony between mechanisms of substrate provision, Ca^{2+} -feedforward and metabolite feedback regulation might underlie acute metabolic inertia. This could be particularly prominent in type II fibres, which have lower CRAT⁸³ and MCU abundance²¹ and slower mitochondrial Ca^{2+} import rates²⁷⁶. Furthermore, metabolic inertia is more pronounced in metabolically compromised and older untrained adults, related to the lower CRAT activity and acetyl-carnitine content of muscle in these individuals⁸⁴.

b, Carbohydrates outcompete non-esterified fatty acids (NEFAs) for oxidation at higher intensities (yellow area). Muscle glucose uptake⁸⁸ and carbohydrate utilization^{71,72,120} increases with exercise intensity. At workloads above maximum fat oxidation (>60–65% of maximal oxygen consumption ($\dot{V}\text{O}_{2\text{max}}$))^{71,72}, flux of pyruvate to acetyl-CoA progressively exceeds rates of TCA cycle entry at citrate synthase (CS), leading to depletion of the muscle free-carnitine ($\text{Carn}_{\text{free}}$) pool through CRAT-dependent acetylation to acetyl-carnitine⁸³. After higher-intensity submaximal exercise, acetylation of the free-carnitine pool is greatest in type I fibres¹⁴². Insufficient free-carnitine availability would inhibit NEFA mitochondrial import at the first step of the carnitine shuttle – that is, carnitine palmitoyl transferase 1B (CPT1B) conjugation of carnitine to long-chain acyl-CoA (LCaCoA). Reduced fat oxidation is associated with diminished free-carnitine levels at ~70% of $\dot{V}\text{O}_{2\text{max}}$ (ref. 72), whereas medium-chain NEFA metabolism bypasses carnitine shuttling and is maintained at higher submaximal workloads²⁷⁷. Therefore, free-carnitine levels appear rate-limiting for long-chain NEFA utilization at increasing exercise intensities. **c**, Lactate and pyruvate oxidation and NADH shuttles (blue area). Downstream of glycolysis, pyruvate (Pyr^-) and/or lactate (La^-) pass through voltage-dependent anion

channels (VDAC), where lactate is converted to pyruvate by mitochondrial lactate dehydrogenase (mLDH) in the intermembrane space¹¹⁹ (step 1). Pyruvate then enters the mitochondrial matrix through the mitochondrial pyruvate carrier (MPC)^{81,119}. The glycerol-3-phosphate (G3P^{2-}) shuttle (G3PS) and malate (Mal^{2-})/aspartate (Asp^{2-}) shuttle enables mitochondrial oxidation of lactate and pyruvate through compartmentalized redox shuttling. G3PS and MAS recycle extra-matrix nicotinamide adenine dinucleotide (NAD^+) (step 2) and transport reducing power from glycolysis to the mitochondrial matrix. This occurs through reactions associated with Mal^{2-} and Asp^{2-} delivery into the matrix space^{81,119} (step 3) and G3P^{2-} donation of electrons directly to coenzyme Q (CoQ) of the electron transport chain¹¹⁹ (step 4). As such, saturation of these shuttles increases lactate accumulation and upregulates the lactate-favouring *LDHA* isoform in vitro²⁷⁸. See Box 3 for discussion of CrP, intermyofibrillar and intramyofibrillar glycogen ($\text{glycogen}_{\text{inter}}$ and $\text{glycogen}_{\text{intra}}$, respectively) and intermyofibrillar lipid droplet (IMF_{LD}) stores, and section 'Acute exercise metabolism in skeletal muscle' for details of their metabolism during acute exercise. β -Ox, β -oxidation; ACSLI, acyl-CoA synthetase long-chain family member 1; AGE, aspartate/glutamate exchanger (also known as SLC25A12); ANT, adenine nucleotide translocator; ATGL, adipose triacylglyceride lipase; *b* → *a* denotes posttranslational modification of PYGM from its less-active *b* form to the constitutively active *a* form by PHK; C, cytochrome c; CACT, carnitine acylcarnitine translocase; mCKM mitochondrial creatine kinase muscle; sCKM, sarcoplasmic CKM; DHAP²⁻, dihydroxyacetone phosphate; DHPR tetrad, four dihydropyridine receptors associated with one ryanodine receptor 1 (RYR1); FABPpm, fatty acid binding protein plasma membrane; FAD, flavin AD; FADH₂, reduced FAD; FATP, FA transporter protein; GLUT4/1, glucose transporters 4 and 1; G-1-P, glucose-1-phosphate; G-6-P, glucose-6-phosphate; HK, hexokinase; HSL, hormone-sensitive lipase; IMS, intermembrane space; LCaCarn, long-chain acyl carnitine; LPL, lipoprotein lipase; sLDH, sarcoplasmic LDH; mMAS, mitochondrial MAS; MCT4/1, monocarboxylate transporters 4 and 1; mG3PDH, mitochondrial glycerol-3-phosphate dehydrogenase; MOE, malate/2-oxoglutarate exchanger (also known as mitochondrial 2-oxoglutarate/malate carrier, M2OM); NADH, reduced NAD; NEFA-alb, albumin-bound NEFA; NOX2, NAD phosphate oxidase 2; O₂⁻, superoxide; 2OG²⁻, 2-oxoglutarate; OXPHOS, oxidative phosphorylation and associated respiratory complexes; PHK, muscle phosphorylase kinase; PYGM, glycogen phosphorylase muscle-associated; ROS, reactive oxygen species; SERCA, sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase; sG3PDH, sarcoplasmic glycerol-3-phosphate dehydrogenase; sMAS, sarcoplasmic MAS; T-tubule, transverse tubule; VLDL1, very low density lipoprotein 1.

enrichment of long-chain fatty acid transporters (fatty acid translocase (FAT, also known as CD36), plasma membrane-associated fatty acid binding protein (FABPpm) and fatty acid transport protein 1 (FATP1) and FATP4)¹³², which have varying capacities for increasing NEFA uptake and oxidation in muscle¹³³. Upon exercise, the exocytosis of CD36 (and possibly other transporters) seems independent of AMPK¹³⁴ but could involve calcium–CaMK kinase (CaMKK)¹³⁵ and MAPK–ERK kinase 1 (MEK1) and MEK2 signalling¹³⁶. Furthermore, although FABPpm is structurally identical to mitochondrial aspartate aminotransferase (mAspAT, also known as AATM), these proteins serve distinct functions within their respective subcellular compartments (that is, FABPpm-mediated transport of long-chain NEFAs across the sarcolemma versus mAspAT-based delivery of reducing equivalents into mitochondria)¹³⁷.

Compared to the situation at rest, NEFAs entering exercising muscle are preferentially directed towards breakdown rather than to re-esterification and storage¹³⁸. Before being used by downstream metabolic pathways, long-chain NEFAs must be activated by thioesterification to long-chain fatty acyl-CoA. Several fatty acid transport proteins possess intrinsic acyl-CoA synthetase activity¹³⁹, but acyl-CoA synthetase

long-chain family member 1 (ACSL1) is the predominant and critical ACSL isoform in muscle¹⁴⁰. As the inner mitochondrial membrane is impermeable to long-chain NEFAs, long-chain fatty acyl-CoA molecules are delivered into the mitochondrial matrix through the 'carnitine shuttle', where they undergo subsequent β -oxidation and OXPHOS.

Prolonged high-intensity exercise resulting in elevated rates of lactate production might directly impede IMCL lipolysis over time by downregulating cAMP–PKA signalling¹⁴¹. However, the glycolytic flux of higher-intensity exercise most probably outcompetes NEFAs for oxidation by depleting the muscle free-carnitine pool^{72,142}. This limits the capacity of the carnitine shuttle and thus impairs the mitochondrial import of long-chain fatty acids (Fig. 2b). Longer exercise durations, lower intramuscular glycogen stores, higher dietary fat intakes and greater type I fibre abundance and aerobic fitness ($\dot{V}\text{O}_{2\text{max}}$) levels can reduce carbohydrate reliance and increase fat utilization during physical activity¹²⁰. Nevertheless, although these factors can delay the transition from predominant fat usage to predominant carbohydrate usage (referred to as the substrate 'crossover point'), the biphasic pattern of NEFA oxidation with exercise intensity^{71,72} is ultimately maintained^{77,143}.

Box 3

Energy depots in skeletal muscle

Creatine phosphate

The total concentration of muscle creatine is ~120–130 mmol per kg dry mass in young adults³²⁰. Approximately 66–67% of this muscle creatine pool is creatine phosphate (CrP), with the rest existing in unphosphorylated form (free creatine)³²⁰. Resting CrP levels are ~10–15% higher in type II fibres, which rely on CrP to a greater extent than type I fibres⁷⁵. CrP is the main ATP substrate of the phosphagen energy system through a reversible reaction catalysed by sarcoplasmic creatine kinase (CKM). At physiological pH during exercise (~6.5–7.0)^{54,55}, CKM is bound to sarcomeres through association with myomesins (MYOM1 and MYOM2) at the M-line³²¹ and phosphofructokinase (PFK) at actin filaments of the I-band³²²—coupling CKM to contraction and glycolysis. CrP hydrolysis peaks at the onset of maximum contraction but deteriorates within ~6 s (ref. 70), and stores can be ~75–90% depleted after roughly 30 s of hard exercise^{70,75}. The majority of CrP resynthesis is linked to aerobic metabolism through the ‘creatine phosphate shuttle’, whereby mitochondrial CKM re-phosphorylates free creatine to CrP in the intermembrane space, using ATP from oxidative phosphorylation³²³ (Fig. 2). Intramuscular creatine rarely reaches saturation by diet and de novo synthesis alone, and supplementing an additional ~3 g per day (~0.03 g per kg body mass daily) increases stores ~15% (to ~140 mmol per kg dry mass) in ~28 days, raising mostly free creatine³²⁴. Creatine supplementation appears safe for renal function³²⁵ and consistently improves strength performance³²⁶. The role of creatine in brain health is an emerging area of interest and creatine supplementation may improve the memory function of older adults³²⁷.

Glycogen

Muscle glycogen concentrations vary depending on nutritional state but are ~400–500 mmol per kg dry mass in the vastus lateralis of individuals following mixed macronutrient diets³²⁸. Glycogen granules are nonuniformly distributed between three specialized pools in muscle³²⁹: (1) the intermyofibrillar pool, found between myofibrils, close to mitochondria and the sarcoplasmic reticulum; (2) the subsarcolemmal pool, positioned just beneath the cell membrane; and (3) the intramyofibrillar pool, situated within the myofibril at the Z-line of the I-band (Fig. 1a). The intermyofibrillar pool is most abundant and accounts for ~77–84% of muscle glycogen, whereas intramyofibrillar and subsarcolemmal stores constitute ~8–15% and ~6–12%, respectively^{35,96,97,267,268}. The relative concentration of each glycogen reserve appears unaffected by age²⁶⁷, type 2 diabetes (T2D)²⁶⁸ or anatomical location (triceps brachii compared to vastus lateralis)³⁵, but may vary depending on physical activity status^{35,267,268} and fibre type^{35,97}. Total glycogen content and utilization is typically higher in type II versus type I fibres^{75,97}, and type II fibres have ~23% more glycogen in intramyofibrillar stores⁹⁷.

Intramyofibrillar glycogen fuels myosin and sodium/potassium (Na⁺/K⁺)-ATPases³²⁹ and is preferentially mobilized during strenuous endurance³⁵, high-intensity interval^{197,330} and resistance⁹⁶ exercise. Myosin-ATPases also utilize intermyofibrillar glycogen, as does

the sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase (SERCA)³²⁹ (Fig. 2). Thus, sufficient muscle glycogen is important for many types of exercise and a critical threshold of ~250 mmol per kg dry mass has been proposed³³⁰, below which self-perceived level of effort increases³³⁰ and sarcoplasmic reticulum function^{330,331}, power output³³¹ and repeated-sprint ability³³⁰ decline until glycogen stores are replenished^{330,331}. Nevertheless, exercising with low glycogen availability (~100–300 mmol per kg dry mass) might augment signal transduction (for example, 5'-AMP-activated protein kinase (AMPK), p53) and gene expression (for example, *PGC1A* and *TFAM*)³³² associated with oxidative metabolism and mitochondrial biogenesis (see the main text). This has led to the concept of ‘train low, compete high’ or ‘carbohydrate periodization’ (reviewed in ref. 332). Whether carbohydrate periodization leads to improved exercise performance over time awaits confirmation, as does delineation of the potential underlying factors (for example, glycogen content per se versus hypocaloric diets that cause weight loss³³³ and/or proximal glycogen-depleting exercise bouts that result in higher training volumes²⁴⁵).

Intramyocellular lipids

Intramyocellular lipids (IMCLs) are stored in the hydrophobic core of lipid droplet ellipsoids⁷⁷ at peripheral (subsarcolemmal, SS_{LD}) and central (intermyofibrillar, IMF_{LD}) regions within fibres^{76–78,268}. Women may have ~43% more individual lipid droplets in muscle, contributing to a greater (~84%) density of total IMCLs than in men³³⁴. Similarly, type I fibres utilize more IMCLs during moderate-intensity exercise¹³⁰ and have ~2–3-fold higher IMCL contents and lipid droplet numbers than type II fibres^{76,78}. Although IMCLs are mostly deposited (>85%) in IMF_{LD}^{76–78}, the relative distribution and characteristics of lipid droplet subpopulations (particularly in type II fibres) vary depending on training status, body composition and metabolic health⁷⁸.

Endurance-trained athletes and adults with T2D have similar total IMCL concentrations in muscle^{78,335} despite markedly dissimilar insulin sensitivity profiles^{78,84,335}. This apparent contradiction has been termed the ‘athlete paradox’³³⁵ but could be partly explained by contrasting lipid droplet properties^{77,78}. Individuals with T2D have a greater number of extremely large SS_{LD} in type II fibres^{77,78}, which also possess lower subsarcolemmal mitochondrial contents⁷⁷. This results in higher relative contributions of SS_{LD} to the overall IMCL pool⁷⁸ and fewer mitochondria-to-SS_{LD} contacts⁷⁷. Spatially, SS_{LD} could interfere with insulin signalling and the T2D-SS_{LD} phenotype negatively correlates with peripheral insulin sensitivity⁷⁸. Conversely, endurance athletes have approximately twofold more IMCLs in type I fibres and an increased abundance of adipose triacylglyceride lipase (ATGL) and perilipin 5 (PLIN5)⁷⁸—proteins associated with lipid droplet turnover¹¹⁸. The higher type I fibre IMCL content of trained individuals is specifically stored in smaller IMF_{LD}⁷⁸, which are favourably depleted during prolonged endurance exercise⁷⁶ (Fig. 2b).

(continued from previous page)

An 8-week programme of high-intensity interval training reduced SS_{LD} size, increased the subsarcolemmal mitochondria-to-lipid droplet ratio and redistributed IMCLs into small IMF_{LD} in type II fibres⁷⁷. Accordingly, lipid droplet profiles were similar between

adults with normal weight, overweight or obesity, or T2D after the training intervention, regardless of baseline differences⁷⁷. Therefore, consistent exercise may alleviate muscle insulin resistance somewhat through remodelling of lipid droplets^{77,78}.

Biological rhythms and skeletal muscle

Normal circadian fluctuations in behaviour and physiology – such as sleep–wake cycles, nutritional state, body temperature, cardiovascular function, and tissue production and sensitivity to hormones (Supplementary Table 1) – coincide to influence acute exercise capacity¹⁴⁴ and response in a time-specific, tissue-specific manner²⁰. Endogenously generated circadian rhythms are predominantly maintained by photic (light-dependent) cues relayed through the central pacemaker (hypothalamic suprachiasmatic nucleus)¹⁴⁵ in concert with cell-autonomous clocks in peripheral tissues, including skeletal muscle (reviewed in ref. 146). Muscle clocks regulate transcriptional programmes that prepare the tissue for transitions between fasting and feeding^{147–149} and orchestrate 24-h rhythms in muscle glucose^{147,149}, lipid^{148,149} and amino acid¹⁴⁸ metabolism. Consequently, muscle clock disruption in mice impairs muscle insulin sensitivity^{147,150} and metabolic homeostasis^{147,148,150}.

Hormonal, metabolic and temperature-dependent changes associated with exercise shift the central pacemaker¹⁵¹ and impact circadian clocks in cell types throughout the body¹⁵². In human vastus lateralis muscle, acute endurance¹⁵³ and resistance^{153,154} exercise upregulates the core clock genes *CRY1* and *PER2*, in part through Ca²⁺-mediated activation of cAMP response element-binding protein (CREB)¹⁵³. Reciprocally, physical activity patterns directly modulate the expression, phase and/or amplitude of -15–20% of all rhythmic genes in mouse soleus and tibialis anterior muscles independently of the muscle clock¹⁵⁵. This occurs through transcription factors such as NFATc1 that are stimulated by motoneuronal firing patterns and downstream Ca²⁺ dynamics¹⁵⁵. Together, these studies provide a mechanistic and biochemical rationale for observed time-dependent variations in human athletic performance (reviewed in ref. 156) and the metabolic disturbances^{157,158} and increased risk of cardiovascular disease¹⁵⁹ resulting from circadian misalignment.

A high-intensity exercise intervention in humans was shown to offset the detrimental impact of short-term sleep restriction on whole-body glucose tolerance and mitochondrial respiration in permeabilized muscle fibres¹⁶⁰. Treadmill running exercise in mice also stimulated hypoxia-inducible factor-1 α (HIF1 α) and a broad range of muscle transcriptomic¹⁶¹ and metabolic^{20,161} responses in a time-dependent manner. Such findings have prompted researchers to begin investigating exercise as a therapy to promote circadian alignment and ameliorate metabolic disease. Preliminary human evidence indicates that training in the afternoon or evening might enhance the beneficial effects of exercise on aspects of insulin sensitivity^{162,163} and blood glucose control¹⁶⁴. Furthermore, moderate-to-vigorous physical activity imparts greater risk reductions for cardiovascular disease and all-cause mortality when $\geq 50\%$ of the total activity volume is undertaken after 11:00 h (ref. 165).

Although rapidly emerging, this branch of ‘chrono-therapeutics’ is still in its infancy, and precise protocols to elicit specific metabolic outcomes are only beginning to come to fruition²⁰. Additionally, there is large population-level variation in human chronotypes¹⁶⁶, and individual chronotype may interact with diurnal exercise timing to modify acute responses and longer-term adaptations. Each age group contains

a normal distribution of early-to-late chronotypes, yet the average chronotype differs according to biological sex (Supplementary Box 2) and changes during ageing¹⁶⁶ (Box 2). Individuals with later chronotypes could benefit from exercise-induced phase advances irrespective of exercise timing, whereas earlier chronotypes might suffer circadian misalignment with evening exercise¹⁵¹. Moving forward, chronotype will be an important covariate to consider in exercise biology, especially for the goal of making personalized therapeutic recommendations.

Skeletal muscle responses to acute exercise

The perturbation of systemic^{19,124} and local⁵⁵ homeostasis posed by acute exercise triggers an integrated series of metabolic, hormonal, growth factor-related, inflammatory and mechanosensitive events. In this section we discuss how these stimuli converge to temporally regulate the acute post-exercise adaptive landscape in muscle – from signal transduction⁹⁰ to gene expression^{167–169} (Fig. 3).

Post-exercise signalling

Sprint and resistance exercise impart stronger whole-body stress responses and more robustly impact the muscle phosphoproteome than continuous endurance exercise does, although there is similarity in the immediate post-exercise signature⁹⁰. Over 400 phosphorylation sites on >200 proteins were commonly altered among these exercise modalities⁹⁰, leading to a shared enrichment of canonical exercise regulatory kinases, including CaMK, AMPK, mammalian target of rapamycin (mTOR), mitogen-activated protein kinase (MAPK), PKA and PKC^{90,170}. Much of this post-exercise pathway convergence is conserved in contracted (exercised) rat and mouse muscle, despite little overlap at specific phosphorylation sites¹⁷¹.

Modality-driven divergence in the phosphoproteome becomes more apparent during the 3-h recovery period after exercise⁹⁰ and may facilitate distinct muscle adaptations with endurance, sprint interval or resistance training. Phosphorylation of rapamycin-sensitive substrates of mTOR complex 1 (mTORC1) and p38 MAPK are potentiated after resistance exercise⁹⁰. mTORC1 is a central regulator of muscle hypertrophy that critically permits mechanical overload-induced myofibrillar accretion¹⁷². In murine muscle, mTORC1 activity is in part mediated by mechanostimulated diacyl glycerol kinase- ζ (DGK ζ) production of phosphatidic acid¹⁷³ and raptor-dependent translocation of the catalytic component of mTORC1 to the late endosomal and lysosomal system¹⁷². By contrast, mTOR in human muscle already associates with the lysosome, and resistance exercise does not further augment this interaction¹⁷⁴. Thus, the contraction-induced migration of the mTOR–lysosomal complex to focal adhesions¹⁷⁵ and microvasculature¹⁷⁴ at the sarcolemma may be a more important step in humans^{174–176}. Here, mTOR associates with RAS homologue enriched in brain (RHEB) and eukaryotic translation initiation factor 3 (EIF3)¹⁷⁴, and activates downstream target ribosomal protein S6 (RPS6)¹⁷⁵. The cooperation between mTOR–EIF3 (ref. 174) and Ser235 and Ser236 phosphorylation of RPS6 (ref. 175) at the fibre periphery was greater when a post-exercise protein and carbohydrate beverage was consumed, and this interplay

Box 4

Select factors produced from skeletal muscle during exercise

Enhanced blood flow during exercise not only improves delivery of nutrients and hormones (Supplementary Table 1) to muscle but also facilitates the release and transport of discrete factors from muscle²⁰. These secreted molecules (known as exercise-induced ‘myokines’ or muscle-derived ‘exerkines’) can act in an autocrine, paracrine or endocrine fashion and are thought to promote many of the favourable adaptations associated with physical activity¹¹.

Interleukin 6 (IL-6) is perhaps the prototypical example of an exercise-stimulated, muscle-secreted factor³³⁶. Elevations in circulating IL-6 with endurance-type exercise^{19,336,337} might contribute towards short-term energy allocation by transiently inhibiting inflammatory processes (for example, monocyte production of tumour necrosis factor (TNF)) while preferentially directing liberated non-esterified fatty acids (NEFAs) towards working muscle (reviewed in ref. 338). Accordingly, systemic pharmacological blockade of the IL-6 receptor promotes re-esterification and storage of NEFAs³³⁷. A ventromedial hypothalamic circuit of locally synthesized IL-6 controls NEFA oxidation after swimming exercise specifically in the soleus of mice, through sympathetic α_{2A}/α_{2C} adrenoceptor modulation of 5'-AMP-activated protein kinase (AMPK)-acetyl-CoA carboxylase (ACC)³³⁹. The effects of centrally produced IL-6 preceded a rise in peripheral IL-6 concentrations, and whether IL-6 from muscle can activate this same neuromuscular axis is unclear³³⁹.

Lactate is an established signalling metabolite and its production is enhanced by glycolytic stressors. Elevated rates of muscle glycolysis during higher-intensity (for example, sprint and resistance) exercise increases plasma lactate to greater extents than moderate-intensity endurance exercise⁹⁰. Muscle-derived lactate can initiate an adipose tissue-transforming growth factor $\beta 2$ (TGF $\beta 2$) secretion axis, which mediates improvements in murine glucose tolerance after 11 days of voluntary wheel running³⁴⁰. Lactate exiting muscle can be further converted to *N*-lactoyl-phenylalanine (Lac-Phe) in cells expressing *CNDP2*, such as immune cell populations (for example, macrophages and monocytes) or epithelial cell populations³⁴¹. Intraperitoneal Lac-Phe delivery caused appetite suppression and

weight loss in obese mice³⁴¹. However, post-exercise concentrations of Lac-Phe in human plasma were orders of magnitude lower than those administered in the murine experimental model³⁴¹. As such, the role of Lac-Phe in the hunger-suppressing effects ~0–3 h after high-intensity exercise³⁴² warrants additional study.

Apelin (APLN)³¹⁵ and succinate³⁴³ are other examples of molecules released from exercising muscle that promote crosstalk between muscle fibres and resident mononuclear cells. This retrograde signalling is particularly important for adaptive extracellular matrix remodelling and angiogenesis (Supplementary Fig. 3). Longer-duration high-intensity intervals⁵⁵ and prolonged endurance exercise¹²⁴ may increase circulating succinate more than resistance exercise does. Similarly, only endurance exercise lowered the [kynurenine (KYN)] to [kynurenic acid (KYNA)] ratio in plasma¹²⁴. KYN is a neurotoxic metabolite and the systemic reduction in [KYN] to [KYNA] protects against stress-induced depressive-like symptoms in mice³⁴⁴. This occurs through a muscle peroxisome proliferator-activated receptor-coactivator 1 α (PGC1 α) isoform 1 (PGC1 α_1)-peroxisome proliferator-activated receptor- α/δ (PPAR α /PPAR δ)-KYN aminotransferase (KAT) cascade that detoxifies KYN to KYNA³⁴⁴. KYNA subsequently released from muscle could influence energy homeostasis by activating G protein-coupled receptor 35 (GPR35) on adipocytes to stimulate lipid turnover³⁴⁵. Numerous other exercise-responsive secretory factors regulated by PGC1 α can reportedly signal from muscle (reviewed in ref. 223), including neurturin (NRTN)³⁴⁶. NRTN operates through both autocrine and paracrine mechanisms to coordinate slow-oxidative muscle fibre and slow-twitch motor neuron property transitions in mice, and its mRNA is upregulated in human vastus lateralis ~72 h after sprint interval exercise³⁴⁶.

Collectively, cell-to-cell and inter-organ communication appears to have an important role in the local and global effects of exercise. Although muscle is a mediator of this crosstalk, other metabolically active tissues (for example, the heart, liver, adipose, nervous, endocrine and immune systems) also contribute, as discussed in detail elsewhere (reviewed in ref. 111).

could contribute towards the nutrient-sensitizing effects of resistance exercise on mTORC1 activation^{6,177}.

Complementary contraction-responsive mechanisms act in tandem with mTORC1 to promote gains in muscle mass^{6,178–180}. For example, sarcomere shortening recruits the β -isoform of MAP3K20 (also known as ZAK β) to the Z-disc, resulting in downstream stimulatory phosphorylation of p38 MAPK and JUN N-terminal kinase 1 (JNK1) and JNK2 (ref. 179). Inhibition mediated by JNK1 and JNK2 (ref. 181) and/or Notch¹⁸²-mediated inhibition of myostatin-suppressor of mothers against decapentaplegic (SMAD) signalling is a purported molecular ‘switch’ in muscle, promoting resistance-type over endurance-type adaptations by preventing SMAD complex nuclear translocation¹⁸¹. This suppression of the myostatin-transforming growth factor- β (TGF β) pathway might be distinct from the regulation of myostatin gene expression¹⁸², as both resistance and endurance training downregulate myostatin mRNA levels in human muscle¹⁸³ despite contrasting hypertrophic phenotypes over time¹³.

Endurance training has been suggested to impair resistance training adaptations through AMPK-mediated blunting of mTORC1. Although this biochemical reaction is known to occur, exercise induction of both AMPK and mTORC1 signalling is consistent with meta-analyses indicating little, if any, interference effect of concurrent exercise on muscle hypertrophy^{184,185}. Bidirectional AMPK-mTORC1 phosphorylation¹⁷⁷ could coordinate simultaneous⁹⁰ or temporally distinct¹⁷⁷ post-exercise activity and downstream substrate specificity of these kinases.

Physiological proteome and mitochondrial remodelling with exercise requires the turnover of select proteins, and autophagic and proteasomal trafficking are integral to these processes. Indeed, genetic disruption of proteostasis results in the accumulation of damaged proteins and impairs force production in fast-twitch extensor digitorum longus and slow-twitch soleus mouse muscle¹⁸⁶. Treadmill running exercise increases the activity of an AMPK complex composed of α_1 , α_2 , β_2 and γ_1 isoforms located on the outer mitochondrial membrane (‘mitoAMPK’)

that could spatially regulate quality control through mitophagy across the mitochondrial reticulum¹⁸⁷. This pool of mitoAMPK is conserved in human muscle¹⁸⁷ and may govern mitophagy through phosphorylation of downstream substrates such as UNC-51-like autophagy activating kinase 1 (ULK1)¹⁸⁸ and mitochondrial fission factor (MFF)¹⁸⁹. In humans, acute high-intensity^{190,191} and sprint interval¹⁹⁰ exercise rapidly and transiently activates ubiquitin-proteasomal degradation in a cAMP–PKA-dependent manner^{190,191} that requires the proteasomal subunit PSD11 (ref. 190). cAMP can then upregulate enzymatic activity of E3 ubiquitin ligases for continued removal of damaged proteins in the post-exercise recovery period¹⁹¹ (Fig. 3). Coordinated proteolytic signalling is further detected 3 h after resistance exercise, as implied by co-regulated phosphorylation of proteins in the N-end rule, calpain regulation and ubiquitin receptor RAD23 pathways⁹⁰.

Many exercise-responsive phospho-sites have yet to be functionally interrogated in muscle^{90,170,178}. Additionally, there are numerous other protein modifications such as arginine methylation, ubiquitin-like modification, ribosylation and acylation that remain relatively understudied in the context of physical activity despite presumably coinciding to influence muscle phenotype. Deeper coverage of the posttranslational proteome combined with the use of subcellularly targeted biosensors¹⁸⁷ should enhance the spatial understanding of how integrated post-exercise signalling governs unique exercise adaptations in muscle, and how these networks change with exercise training over time.

Epigenetic regulation of exercise adaptation

Histone modifications and DNA methylation. Upstream signalling, together with an altered intramuscular metabolic milieu, can change the chromatin landscape in muscle and thus influence transcription factor-driven gene expression in response to exercise (reviewed in ref. 192). Epigenetic modifications on amino acids in histone tails or globular core domains render a permissive or repressive chromatin state depending on the type of modification and the residue and/or histone protein being targeted¹⁹². An acute bout of endurance exercise can increase H3K36 acetylation¹⁹³, a mark of euchromatin, and promote CaMKII-dependent phosphorylation^{194,195} and nuclear exclusion of the transcriptionally repressive class II histone deacetylases (HDAC) HDAC4 and HDAC5 (ref. 193). High-intensity resistance exercise also transiently stimulates myonuclear H3S10 phosphorylation through p38 MAPK and mitogen and stress-activated protein kinase 1 (MSK1) in human muscle¹⁹⁶. This same axis was necessary for the upregulation of myocardin-related transcription factor B (MRTFB)–serum response factor (SRF)-target genes and protein synthesis after *in vivo* contractions in mice¹⁹⁶.


Post-exercise histone modifications coalesce with changes in DNA methylation, typically at the 5' end of cytosine residues at CpG sites throughout the genome. DNA methylation in *cis*-acting regulatory regions is commonly associated with gene silencing, but some transcription factors, particularly of the homeodomain POU and NFAT families, favour methylated CpG sequences¹⁹⁷. Acute exercise^{198–200} and exercise training^{199,201} induce both hypomethylation and hypermethylation of DNA in muscle, mostly within gene bodies, intergenic regions and enhancer regions²⁰¹. The greatest number of CpG-site methylation changes and the highest percentage of mRNA transcripts with inversed DNA methylation patterns occurred 3 h after a single session of resistance exercise, irrespective of workload (80% or 30% of 1RM)²⁰⁰. These changes began to revert towards baseline methylation levels by 6 h²⁰⁰, suggesting that the muscle methylome is dynamic and tightly regulated during recovery from exercise. Yet, select DNA hypomethylation and

hypermethylation events are sustained after acute resistance exercise and resistance training, providing a plausible mechanism for a 'muscle memory' of previous exercise that facilitates future adaptation¹⁹⁹. Modality-dependent DNA methylation signatures might also direct specific post-exercise transcriptional programmes in muscle, as acute bouts of resistance¹⁹⁹ and sprinting-type²⁰² exercise share enrichment for only 36 of the top 100 differentially methylated pathways.

Conceivably, exercise-induced metabolic fluxes, which stimulate glycolysis and OXPHOS, could alter acetyl and methyl group availability, cellular [NAD⁺] to [NADH] redox state and the activity of 'writers' and 'erasers' of epigenetic marks in muscle¹⁹². However, the current understanding of exercise-regulated DNA methylation and chromatin remodelling is incomplete. For example, lactate modifies ('lactylates') H3K18 in promoters and enhancers of genes related to tissue development and metabolism in human muscle²⁰³, but whether lactylation of histone lysine residues contributes to exercise adaptation remains to be ascertained.

Non-coding RNAs. In addition to protein-coding genes, RNA polymerase II transcriptional activity produces microRNAs (miRNAs; ≤ 22 nucleotides) and long non-coding RNAs (lncRNAs; ≥ 200 nucleotides) that alter mRNA levels and protein abundance without influencing the genetic code. Many miRNAs²⁰⁴ and lncRNAs²⁰⁵ are exercise-responsive in muscle, and both classes seem involved in aspects of myogenesis. During resistance exercise, the downregulation of the muscle-enriched miRNA myomiR-1, through its contraction-mediated release in extracellular vesicles, may facilitate hypertrophy²⁰⁶. Once in circulation, myomiR-1 could act upon adipocytes to potentiate adrenergic signalling and lipolysis²⁰⁶. Still, muscle was not a major source of bloodborne extracellular vesicles upon resistance exercise in humans²⁰⁷ or treadmill running exercise in mice²⁰⁸. Rather, muscle-derived extracellular vesicles – loaded with myomiRs – preferentially accumulated in the interstitial space, and treatment with these vesicles promoted differentiation of myoblasts *in vitro*²⁰⁸. Therefore, extracellular vesicles originating from muscle might play a more prominent role in local cellular communication (Supplementary Fig. 3). Satellite cells may signal to muscle fibres²⁰⁹ and fibroadipogenic progenitor cells²¹⁰ using extracellular vesicles containing myomiR-206. In mice, this was shown to regulate the levels of extracellular matrix (ECM)-related genes (for example, *Mmp9*) in muscle fibres²⁰⁹ and *Rrbp1* in fibroadipogenic progenitor cells²¹⁰ to coordinate physiological ECM remodelling during the early stages of hypertrophy. However, the ability of extracellular vesicles to deliver biologically active miRNA cargo into recipient cells remains contentious²¹¹ and their role in exercise adaptation warrants further interrogation.

Though the impact of lncRNAs in human muscle is less studied than that of miRNAs, lncRNAs respond to endurance, resistance, concurrent and high-intensity interval training with distinct differential expression profiles between modalities²⁰⁵. The lncRNA *CYTOR* is induced by endurance and resistance exercise in human muscle and can regulate fast-twitch fibre formation in rodents by sequestering TEA domain family member 1 (TEAD1) transcriptional activity²¹². *Cytor* mRNA is decreased upon ageing and overexpression of this lncRNA increased type IIA and type IIB fibre abundance, which was sufficient to improve grip strength and uphill running performance in aged mice²¹². RNAs putatively annotated as lncRNAs may also contain short open reading frames that encode for micropeptides (≤ 150 amino acids). The muscle-specific micropeptides dwarf open reading frame²¹³ and myoregulin²¹⁴ competitively interact with sarcoplasmic/endoplasmic



reticulum Ca^{2+} -ATPase (SERCA) to positively and negatively affect SR Ca^{2+} kinetics and muscle contractility, respectively. Other classes of non-coding RNAs, including circular RNAs and tRNA fragments, await future study in the context of human muscle exercise adaptation.

mRNA transport and myonuclear propagation

The dense myofibrillar lattice in adult muscle fibres may impede diffusion of mRNAs expressed from myonuclei. Indeed, recent evidence suggests that mRNA distribution away from the perinuclear region is almost entirely dependent on microtubule networks in muscle, whereas diffusion has a minor role^{215,216}. Transport might be reliant on motor proteins, reminiscent of cardiomyocytes that use kinesin-1-mediated

transport of mRNA and ribosomes to direct protein translation to specific sites within the cell²¹⁷. Microtubule-based trafficking of mRNA facilitates transcript distribution through the crowded sarcoplasmic space, but dispersion of mRNAs from progenitor myonuclei is still potentially confined to $\leq 50 \mu\text{m}$ (refs. 215,216). By contrast, the movement of proteins translated from transcripts derived from a specific myonucleus could be ≥ 5 times more far-reaching²¹⁸.

The use of fluorescent reporters has shown that nuclear proteins can be detected in surrounding myonuclei and in those distant from the nucleus of origin^{218–220}, a phenomenon termed ‘nuclear propagation’²¹⁸. Smaller proteins ($\leq 40 \text{kDa}$) that freely diffuse across the nuclear pore complex can propagate to many myonuclei throughout

Fig. 3 | Molecular responses to acute exercise and exercise training. Exercise-induced alterations in circulating molecules^{19,124} and the intramuscular milieu^{55,101}, together with mechanical tension¹⁷⁸, initiates a temporal series of biochemical and molecular events that lead to muscle adaptation. Activation of signalling cascades promote substantial posttranslational modification of the muscle proteome^{90,170} and DNA accessibility^{198,199,230}. Collectively, this drives transcription factor-dependent¹⁶⁹ changes in gene expression^{167,183}, alongside microRNAs³⁰⁴ and long-non-coding RNAs²⁰⁵ that are thought to ‘fine-tune’ the molecular responses to exercise. Endurance exercise (EE) and resistance exercise (RE) are often considered divergent stimuli, primarily driving oxidative versus hypertrophic muscle adaptations, respectively. However, common processes among exercise modalities can result in shared enrichment of signalling cascades⁹⁰ and transcriptional networks¹⁸³ in the post-exercise period. For example, coordinated proteolysis is detected following acute exercise, irrespective of exercise type⁹⁰. This may require cAMP–protein kinase A (PKA)^{190,191} and ensures protein quality control and physiological muscle remodelling. 5′-AMP-activated protein kinase (AMPK) activity⁹⁰ and the expression of total *PGC1α* mRNA¹⁸³ are also increased after a single bout of either endurance or resistance exercise. AMPK phosphorylation activates peroxisome proliferator-activated receptor-γ coactivator 1α isoform 1 (*PGC1α*)²⁷⁹, and both AMPK and *PGC1α* potentiate angiogenic factors^{260,280} and mitochondrial bioenergetics^{48,235,260,279} in muscle. After endurance exercise, a distinct pool of mitochondrial AMPK (composed of α, α₂, β₂ and γ₁ isoforms) regulates mitophagy^{187–189} and promoter hypomethylation facilitates the transcription of peroxisome proliferator-activated receptor-δ (*PPARδ*) and transcription factor A, mitochondrial (*TFAM*)²³⁰. Whether resistance exercise elicits these same effects is unclear. Despite similarities, there are more distinct than overlapping post-exercise responses between modalities^{90,183}. Rapamycin-sensitive substrates of mammalian target of rapamycin (mTOR) complex 1 (mTORC1) are phosphorylated to a greater extent after resistance exercise⁹⁰. Mechanical overload initiates translocation of the mTOR–lysosomal complex¹⁷⁴ and diacylglycerol (DAG) kinase-ζ (*DGKζ*)¹⁷³ to the sarcolemma. Here, mTOR colocalization with RAS homologue enriched in brain (*RHEB*) and eukaryotic initiation factor 3 (*EIF3*)¹⁷⁴ and phosphatidic acid (PA) produced by *DGKζ*¹⁷³ might coalesce to fully stimulate mTORC1 (ref. 175) and the translation of contraction-associated mRNAs. Acute attenuation of UNC-51-like autophagy activating kinase 1 (*ULK1*) autophagic signalling after resistance exercise⁶⁵ and nuclear *DGKζ*-mediated suppression of forkhead box protein O (*FOXO*)-dependent proteasomal degradation could also support muscle mass by moderating the breakdown of myofibrillar proteins¹⁷³. At the sarcomere, contraction recruits ZAKβ to the Z-disc¹⁷⁹ where it acts through JUN N-terminal kinase 1 (*JNK1*) and *JNK2* (refs. 179,181), potentially alongside Notch¹⁸², to inhibit

myostatin (*MSTN*)/transforming growth factor-β (*TGFβ*) signalling. This represents one of many intracellular changes permitting resistance training-induced hypertrophy over endurance-like adaptations¹⁸¹. The upregulation of *MYC* with resistance exercise stimulates ribosomal biogenesis^{180,198}, and the formation of a specialized pool of ribosomes with a high ratio of ribosomal protein large 3 (*RPL3*) to *RPL3*-like (*RPL3L*)²³⁸ may favour protein synthesis over translational fidelity. *MYC* expression is mTOR-independent but *MYC* cooperation with mTOR is necessary to successfully increase ribosomal content¹⁸⁰, possibly requiring an mTOR-driven reorganization of nucleoli to aid rRNA transcription²⁸¹. Divergence between endurance and resistance exercise at the transcriptomic level is magnified after a period of training¹⁸³. Endurance training increases electron transport chain complex expression¹⁸³, mitochondrial content and muscle oxidative capacity¹². Conversely, growth-related pathways¹⁸³, ribosomal abundance¹⁹⁸ and muscle mass¹² are augmented more by resistance training. This could be mediated in part by *PGC1α* isoforms. Nuclear localization of *PGC1α*, and Ser15 phosphorylation of p53 are greater in resting muscle after high-intensity interval training²³⁹, which might help to preserve mitochondrial content and function²⁵⁷. By contrast, *PGC1α* protein is unchanged after resistance training, whereas the *PGC1α* isoform 4 (*PGC1α₄*) protein is preferentially enriched²³⁵. *PGC1α₄* stimulates muscle hypertrophy and is associated with enhanced *Igf1* expression²²⁹, insulin-like growth factor 1 (*IGF1*)–serine/threonine protein kinase (*AKT*) and mTORC1 signalling²³⁴ and downregulation of *Mstn* mRNA in mouse muscle²²⁹. However, unlike *PGC1α₁* (ref. 280), *PGC1α₄* does not coactivate oestrogen-related receptor-α (*ERRα*)²²⁹ and has no effect on oxidative phosphorylation enzymes^{229,235}. Appreciable overlap in the initial stages of exercise training probably underlies the degree of shared adaptation between endurance and resistance exercise (see the sections ‘Skeletal muscle responses to acute exercise’ and ‘Skeletal muscle adaptations to long-term exercise’). Depending on individual predisposition (Box 1), dedicated training of a certain exercise modality could amplify discrete differences in the adaptive response, resulting in distinct muscle adaptations and the development of specific phenotypes over time¹³. Evidence showing that combined endurance and resistance training can blunt muscle hypertrophy in humans is scarce^{184,185} but concurrent training could impede gains in explosive strength¹⁸⁴. Still, a combined exercise regime may offer dual benefits for most individuals¹². *ACTRIIB*, activin receptor type 2B; *ECM*, extracellular matrix; *MFF*, mitochondrial fission factor; *MCU*, mitochondrial calcium uniporter; *NFATc1*, nuclear factor of activated T cells, cytoplasmic 1; *NICD*, Notch intracellular domain; *NRF2*, nuclear factor erythroid 2-related factor 2; *p38*, p38 mitogen-activated protein kinase; *RPS6*, ribosomal protein S6; *SMAD*, mothers against decapentaplegic homologue; *Ub*, ubiquitin; *ZAKβ*, *MAP3K20* isoform-β.

a myotube *in vitro*²¹⁸. Conversely, larger proteins (>40–60 kDa) are limited in their propagation potential, possibly due to increasing reliance on nuclear transport receptors to traverse the nuclear pore complex²¹⁸. Protein properties (such as size) and myonuclear characteristics (such as shape or location within the muscle fibre) probably interact to control nuclear propagation. For example, myonuclei in *ex vivo* fibres differ in capacity for classical versus non-classical nuclear import, resulting in a gradient of import rates between myonuclei of the same fibre²¹⁹.

The potential for myonuclear propagation casts further doubt on strict myonuclear domains in muscle and represents an additional layer of spatiotemporal control, combining fibre-wide communication with myonuclei-autonomous processes in response to adaptive cues. Furthermore, resident myonuclei may be capable of endoreplication in mouse muscle (resulting in polyploidy), which could support efficient DNA synthesis under heightened transcriptional demand²²¹. Myonuclear propagation was detected after wheel-running exercise training²²⁰ and synergistic ablation-induced hypertrophy increased

the total number (from -0.7–1% to -3%) of DNA-replicating myonuclei²²¹ in murine muscle. The extent to which these processes occur under physiological conditions in humans remains unclear.

The post-exercise transcriptome

Considerable effort has been dedicated towards characterizing the transcriptional response of human muscle to exercise, and several key studies are highlighted in Supplementary Table 2. High-powered meta-analyses have curated much of this scientific endeavour to generate deeper insight into exercise adaptation^{167,183}. Acute exercise initiates temporal clusters of gene expression^{167–169}, including early induction of stress response genes, increased expression of muscle adaptation genes in early and middle stages following exercise^{167,169}, and later¹⁶⁷ or sustained¹⁶⁸ increases in immune-related signatures. These transcriptional programmes are driven by combinations of different transcription factors^{167,169} and alternative transcription start-site usage¹⁶⁹. Akin to the phosphoproteome⁹⁰, a single bout of resistance exercise alters more transcripts than does moderate-intensity endurance exercise¹⁸³.

However, there is considerable overlap in pathway enrichments, and both training modalities upregulate stress response genes, kinase activity and metabolism-related processes, among others¹⁸³.

NR4A3 (also known as *NORI*)²²² and *PGCIA*²²³ are genes commonly induced after endurance and resistance exercise¹⁸³ and mainly promote oxidative muscle adaptations. *NR4A3* and *PGCIA* have similar temporal expression patterns after exercise, peaking -2–3 h into the recovery period^{167,183}, suggesting common upstream regulation through CREB^{224,225}. The nuclear receptor subfamily 4 group A (NR4A) family (including NR4A1 (also known as NUR77) and NR4A2 (also known as NURR1)) share >90% homology in their DNA-binding domains and can interact as monomers with a single NGFI-B response element²²⁶, or as homodimers and heterodimers (alongside other NR4A members) with the Nur response element²²⁷. *NR4A1* is also exercise-responsive in muscle^{167,202} and may serve partial redundancy with NR4A3, as implied by the slow-oxidative phenotypes of mice expressing either *Nr4a1* (ref. 228) or *Nr4a3* (ref. 222) from skeletal muscle-specific promoters.

The PGC1 α family of coactivators consists of different isoforms, which are derived from alternative promoter use or splicing events (reviewed in ref. 223). Endurance exercise preferentially targets²²⁹ and hypomethylates²³⁰ the proximal PGC1 α promoter in muscle, increasing PGC1 α isoform 1 (PGC1 α_1) expression. PGC1 α_1 drives mitochondrial biogenesis and oxidative fibre properties⁴⁸ at least in part through C-terminal domain RNA binding, which facilitates recruitment in transcriptionally active chromatin condensates²³¹. A bout of endurance exercise also leads to promoter region hypomethylation of other key mitochondrial and metabolic genes, including *TFAM* and *PPARD*, with corresponding increases in mRNA expression²³⁰. The PPAR δ -mediated upregulation of lipid metabolism and suppression of glycolytic processes in muscle²³² is largely dependent on protein–protein interaction with Krüppel-like factor 15 (KLF15)²³³.

Most PGC1 α isoforms have overlapping roles in muscle²²³. The truncated variant PGC1 α isoform 4 (PGC1 α_4) differs in that it can be induced by intramitochondrial [Ca²⁺]²³⁴ and is transcribed from the alternative PGC1 α promoter after resistance exercise^{229,235} (Fig. 3). PGC1 α_4 facilitates muscle hypertrophy^{229,234,235} and a PPAR β –AMPK-dependent upregulation of glycolytic metabolism²³⁵, which could enhance the anaerobic capacity of muscle after resistance training. Incidentally, the accretion of muscle mass often coincides with a glycolytic switch that may support anabolism by increasing de novo synthesis of nucleotides and non-essential amino acids from intermediates in glucose metabolism (reviewed in ref. 236).

An acute bout of resistance exercise also stimulates *MYC* mRNA expression and the hypomethylation of enhancer and MYC-accessible intergenic regions of ribosomal DNA¹⁹⁸. These events coalesce to augment 45S pre-rRNA transcription and ribosomal biogenesis^{180,198}. Whereas mTORC1 signalling promotes translational efficiency^{172,180} and directs ribosomal function through Ser235 and Ser236 phosphorylation of RPS6 (ref. 237), total ribosomal content¹⁸⁰ and the altered composition of ribosomal proteins (for example, an increased ratio of RPL3 to RPL3L)²³⁸ determines translational capacity in muscle. Collectively, upregulation of the sarcoplasmic and mitochondrial protein synthetic machinery (such as mito-ribosomes) enables the beneficial proteome remodelling that occurs with high-intensity interval, resistance or combined exercise training¹².

Skeletal muscle adaptations to long-term exercise

In this section, we contextualize the impact of consistent exercise (exercise training) on muscle phenotype, considering general and

select adaptations to specific training interventions and the influence of muscle fibre type¹⁴. We then summarize how muscle-related adaptations can contribute to improvements in systemic measures of exercise performance and discuss whether altered molecular perturbations in muscle with exercise training^{33,239–241} reflect an attenuated or refined adaptive state.

Transcriptome, proteome and contractile property adaptations

Repeated exposure to specific exercise modalities results in shared and divergent alterations in muscle over time^{12,183} (Fig. 3). At the transcriptomic level, pathways related to ECM remodelling, angiogenesis and the TCA cycle are enriched in muscle after exercise training¹⁶⁷. Whereas resistance exercise robustly effects ECM reorganization^{183,242} and growth-related pathways²⁴², endurance training favourably impacts oxidative metabolic processes and the gene expression of mitochondrial complex subunits¹⁸³. Crosstalk between the various cell populations in muscle appears particularly important for exercise-associated adaptations within the interstitial space (Supplementary Fig. 3).

Training-induced improvements in muscle respiration often coincide with expansion and modification of the mitochondrial proteome^{12,15,143} in type I and type II fibres¹⁴. Resistance training can augment the respiratory and ATP-producing capacity of mitochondria²⁴³, but the effects on mitochondrial phenotype are typically less pronounced than those observed after high-intensity interval^{12,15,143} and endurance¹⁴ training. The total volume of endurance exercise performed during a training intervention might be the strongest stimulus for mitochondrial biogenesis in muscle (reviewed in ref. 244). This could be related to the additive effects of proximal exercise bouts on the nuclear enrichment of transcriptional regulators of mitochondrial abundance (for example, PGC1 α_1 and transcription factor EB (TFEB))²⁴⁵ and the upregulation of mito-ribosomes^{12,14,15} and mitochondrial protein synthesis¹². During mitochondrial biogenesis, stoichiometry of the ETC is maintained through proportional translation of ETC complex subunits from the mitochondrial and nuclear genomes²⁴⁶. Increased expression of mitochondrial transit peptides and translocase proteins of the outer and inner mitochondrial membrane following endurance training¹⁴ facilitates the transport of nuclear-encoded precursor components into the mitochondrial matrix. However, in the early stages of mitochondrial remodelling, synthesis of fatty acid oxidation and TCA cycle enzymes may be prioritized over ETC machinery biogenesis, to enhance reducing equivalent NADH and FADH₂ production and electron delivery to OXPHOS¹⁵.

High-intensity interval training at $\geq 90\%$ of maximum power output (\dot{W}_{\max}) most consistently increases normalized (mass-specific) mitochondrial respiration²⁴⁴, with intensities $\geq 100\% \dot{W}_{\max}$ (as in sprint interval training) doing so more efficiently per unit of time spent exercising²⁴⁴. The impact of intensity on mitochondrial respiration appears uncoupled from total training volume, suggesting that these variables drive complementary but somewhat distinct mitochondrial adaptations²⁴⁴. High-intensity interval training also induces lysine acetylation of TCA cycle and ETC proteins¹⁴³, although whether such posttranslational modifications contribute towards intensity-dependent changes in mitochondrial function requires additional study.

After endurance training, some discrete differences can be seen in the adaptive response of type I and type II fibres¹⁴. Still, most of the commonly detected exercise-regulated proteins related to mitochondrial and glucose metabolism behaved similarly between fibre types¹⁴. For example, training upregulated ACSL1, malate/aspartate shuttle

proteins, the mitochondrial CKM isoform CKMT2, PDH, CRAT and LDHB in both type I and type II fibres¹⁴. *LDHB* mRNA is increased after endurance²⁴⁷ and resistance²³⁵ training, potentially through PGC1 α 1 coactivation of distal myocyte enhancer factor 2 (MEF2) and proximal oestrogen-related receptor (ERR) binding sites in the *LDHB* promoter²⁴⁷. LDHB preferentially catalyses the conversion of lactate to pyruvate and, together with enhanced mitochondrial density, could improve lactate clearance in trained muscle through higher rates of conversion to pyruvate and subsequent oxidation²⁴⁷ (Fig. 2c). Further refinement of substrate handling with programmed exercise is implied by greater expression of key proteins of metabolic pathways involving glycogen (for example, glycogen synthase)¹¹, NAD⁺ (nicotinamide phosphoribosyltransferase) and branched-chain amino acid (branched-chain α -ketoacid dehydrogenase kinase) metabolism, as well as ubiquinone biosynthesis (5-demethoxyubiquinone hydroxylase)¹⁴³, after high-intensity interval training.

Muscle contractile properties are likewise affected by regular exercise in a somewhat training modality-dependent manner. Protein isoforms regulating SR Ca²⁺ release¹⁴³ and myofibrillar Ca²⁺ sensitivity^{14,143} were altered after moderate-intensity endurance¹⁴ and high-intensity interval¹⁴³ training, coincident with a reduction in fibre size-adjusted quadriceps peak-twitch torque and prolonged half-relaxation time¹¹ – indicative of a slower muscle phenotype. Conversely, stretch-shortening cycle (ballistics or plyometrics) training increased contractile velocity, force production and thus peak power of single type I, type IIA and type IIA/IIX fibres⁴². Exercise modalities commonly decrease *MYHI* mRNA¹⁸³ and protein (MyHC-IIX)^{11,13} content in muscle. However, the upregulation of *MYH7* gene expression¹⁸³ and type I fibre abundance after endurance training⁴⁴ can be in contrast to the effects of resistance training, which often downregulates *MYH7* (ref. 183) with no corresponding change in the prevalence of type I fibres⁴⁵. This is consistent with endurance⁴⁴ and resistance⁴⁵ exercise differentially shifting hybrid fibres towards pure type I and pure type IIA fibres, respectively (reviewed in ref. 38). Indeed, endurance training resulted in the modification of more proteins in type I than in type II fibres¹⁴, and long-term exposure to a particular exercise stimulus⁴¹ may largely explain the majority of type I or type II fibres detected in biopsies from endurance or strength athletes¹³.

Training-induced adaptations impact exercise performance

Together, exercise training adaptations in muscle contribute towards improvements in gross performance measures such as $\dot{V}O_{2\max}$ (refs. 12,14,15), peak power output^{11,143}, ballistic power⁴² and strength^{12,13}. Endurance and resistance training increase muscle capillarization^{248–250} and endurance-type training enhances NEFA oxidation at a given exercise intensity^{77,120,143,251}. These glycogen-sparing and metabolite-buffering (for example, lactate clearing)²⁴⁷ effects render muscle more robust to disturbances in metabolic homeostasis, delaying mechanisms of peripheral and central fatigue⁹⁴ and raising exercise tolerance and capacity (that is, higher velocities and percentages of $\dot{V}O_{2\max}$ produced at ventilatory thresholds, and greater maximal running speeds or distances covered)²⁵¹. Metabolomics analysis also implies that 4 weeks (13 bouts) of resistance exercise may suffice to modify the acute muscle metabolome²⁵². Yet, the specific metabolic changes that occur with resistance training are not extensively characterized and warrant further interrogation.

High-intensity interval^{11,12} and endurance¹³ training both induce a degree of muscle hypertrophy, but gains in muscle mass and strength are less pronounced than seen with long-term resistance exercise¹³.

Conversely, any beneficial effect of resistance training on $\dot{V}O_{2\max}$ appears limited to individuals with lower initial aerobic fitness levels (for example, a $\dot{V}O_{2\max} \leq 25$ or ≤ 40 mL kg⁻¹ min⁻¹ in adults >60 and ~20–35 years of age, respectively) (reviewed in ref. 253) and thus high-intensity interval¹² and endurance training^{13,248} typically increase $\dot{V}O_{2\max}$ to greater extents.

A period of aerobic pre-conditioning can potentiate the hypertrophic effects of resistance exercise²⁴⁸, in part by enhancing fibre capillary density^{248,250} (Supplementary Fig. 3). Alternatively, high-intensity resistance training ($\geq 90\%$ of 1 RM or ≤ 4 RM) can improve running economy (that is, reduce the metabolic or $\dot{V}O_2$ cost of running)²⁵⁴. These complementary adaptations further emphasize the utility of incorporating numerous exercise (sub)modalities into a sustainable routine for maximum performance¹² and protective health benefits¹⁸.

Adaptive responses in muscle are modified with exercise training

The recurring stimulus of exercise training dampens select signalling^{239,240} and transcriptomic^{33,241} responses to an acute bout of physical activity. This is evident after just nine sessions of high-intensity interval exercise through a modest reduction in the number (-17%) and amplitude (-30%) of altered mRNA transcripts in human muscle compared to the first exercise bout, including glycolysis pathway and HIF1 α target genes²⁴¹. The magnitude of change in the muscle transcriptome with exercise training becomes even clearer over 30 daily sessions of electrically induced mechanical overload in rat hindlimb muscle³³. In particular, the directionality of altered genes contrasted substantially between exercise naive and trained muscle. At the 1-h sampling time point after exercise, ~70% of the (~2400) differentially expressed transcripts were upregulated on day 2 compared with ~83% (of ~3300 genes) downregulated on day 30 (ref. 33). This indicates that certain signalling mechanisms are sensitive to the modified intracellular environment after a period of regular exercise. In combination with genetic predisposition (Box 1), blunted post-exercise molecular responses probably converge to limit the adaptive potential of muscle, slowing progress with advanced training experience. Nevertheless, some alterations might reflect a refined (as opposed to impaired) response.

Pathways related to ribosomal biogenesis and protein synthesis are positively enriched in the early adaptive stages in exercised rat muscle³³ but become downregulated in favour of metabolism-related processes once hypertrophy plateaus³³. Rates of myofibrillar protein synthesis also correlate better with measures of muscle mass accrual after 8 weeks¹⁷⁶ to 10 weeks²⁵⁵ of resistance training in humans, corresponding to the attenuation of exercise-induced muscle damage²⁵⁵. Furthermore, although 20 days (40 sessions) of high-frequency, high-intensity interval training reduced the post-exercise upregulation of nuclear PGC1 α 1 and p53 phosphorylation at Ser15, resting levels of these same markers were greater in human muscle after the exercise intervention²³⁹. Phosphorylated p53 can translocate to the mitochondrial genome where it interacts with transcription factor A, mitochondrial (TFAM) to regulate mitochondrial transcription²⁵⁶ (Fig. 3). Muscle-specific *Trp53*-knockout mice have reduced basal mitochondrial contents and enzymatic activities but retain the capacity for exercise-stimulated mitochondrial biogenesis²⁵⁷. This suggests that p53 may be more important for the maintenance of muscle mitochondrial integrity than exercise adaptation.

Given these diverse modifications, longer-duration trials are needed to fully elucidate the modality-dependent temporal landscape of exercise adaptation in human muscle and to highlight specific changes in the acute exercise response with consistent training.

Future directions

The complex and intricate macrostructure to ultrastructure of skeletal muscle allows for a high degree of metabolic flexibility during exercise. Further study of the regulation and interaction of key cellular components, such as myofibrillar and mitochondrial networks⁵³, should enable more comprehensive understanding of how muscles mount such a malleable response. The use of fluorescent metabolite²⁵⁸ and myofilament²⁵⁹ biosensors could aid this effort by elucidating real-time dynamics of muscle metabolism and contraction.

Current knowledge of exercise adaptation is heavily biased towards canonical regulators such as AMPK, PGC1s and mTORC1. Although these factors are integral to muscle remodelling with exercise, aspects of adaptation are maintained in the absence of AMPK α_1 and AMPK α_2 (ref. 260) and of PGC1 α and PGC1 β (ref. 261), and rapamycin-insensitive pathways also contribute towards muscle hypertrophy^{178,180}. Most of the exercise-induced phosphoproteome remains relatively unexplored^{90,170,178}, and its functional interrogation will help to identify redundancies in adaptive signalling and new players in the distinct and complementary effects of resistance, endurance and high-intensity exercise.

As exercise biology moves increasingly towards systems-level profiling, the multiplexing of new technologies could provide invaluable insight into the spatiotemporal control of gene expression²⁶², translation²⁶³ and metabolism³⁷ in discrete muscle cell populations. If performed in tandem with commensurate functional assays, this would enable causal association of specific cellular events with their phenotypic impact.

Exercise is an effective adjunct therapy for certain neurodegenerative²⁶⁴ and mental health²⁶⁵ disorders and offers protection against cardiometabolic disease²⁶⁶, sarcopenia (Box 2) and all-cause mortality¹⁸. Multicentre interventions including spectrums of chronological age, metabolic health, chronotype, ethnicity, biological sex and social gender will move the field closer to uncovering the dynamic changes that coordinate the benefits of exercise training and further define the role of muscle in the integrative exercise response. Knowledge derived from such approaches should not only inform personalized exercise prescription, but also reveal new molecular avenues of drug discovery for the improvement of human health.

Published online: 24 May 2023

References

- Bilet, L. et al. One-leg inactivity induces a reduction in mitochondrial oxidative capacity, intramyocellular lipid accumulation and reduced insulin signalling upon lipid infusion: a human study with unilateral limb suspension. *Diabetologia* **63**, 1211–1222 (2020).
- Bergouignan, A. et al. Effect of contrasted levels of habitual physical activity on metabolic flexibility. *J. Appl. Physiol.* **114**, 371–379 (2013).
- Kivimaki, M. et al. Physical inactivity, cardiometabolic disease, and risk of dementia: an individual-participant meta-analysis. *Brit. Med. J.* **365**, 11495 (2019).
- Zhang, D. et al. Leisure-time physical activity and incident metabolic syndrome: a systematic review and dose-response meta-analysis of cohort studies. *Metabolism* **75**, 36–44 (2017).
- Millard, L. A. C., Tilling, K., Gaunt, T. R., Carslake, D. & Lawlor, D. A. Association of physical activity intensity and bout length with mortality: an observational study of 79,503 UK Biobank participants. *PLoS Med.* **18**, e1003757 (2021).
- D'Hulst, G., Masschelein, E. & De Bock, K. Resistance exercise enhances long-term mTORC1 sensitivity to leucine. *Mol. Metab.* **66**, 101615 (2022).
- Goodyear, L. J. et al. Skeletal muscle plasma membrane glucose transport and glucose transporters after exercise. *J. Appl. Physiol.* **68**, 193–198 (1990).
- Burd, N. A. et al. Enhanced amino acid sensitivity of myofibrillar protein synthesis persists for up to 24 h after resistance exercise in young men. *J. Nutr.* **141**, 568–573 (2011).
- Sjoberg, K. A. et al. Exercise increases human skeletal muscle insulin sensitivity via coordinated increases in microvascular perfusion and molecular signaling. *Diabetes* **66**, 1501–1510 (2017).

- McConell, G. K. et al. Insulin-induced membrane permeability to glucose in human muscles at rest and following exercise. *J. Physiol.* **598**, 303–315 (2020).
- Hostrup, M., Onslev, J., Jacobson, G. A., Wilson, R. & Bangsbo, J. Chronic β_2 -adrenoceptor agonist treatment alters muscle proteome and functional adaptations induced by high intensity training in young men. *J. Physiol.* **596**, 231–252 (2018).
- Robinson, M. M. et al. Enhanced protein translation underlies improved metabolic and physical adaptations to different exercise training modes in young and old humans. *Cell Metab.* **25**, 581–592 (2017).
This study provides a thorough analysis of exercise adaptation at several -omics levels in human skeletal muscle.
- Chapman, M. A. et al. Skeletal muscle transcriptomic comparison between long-term trained and untrained men and women. *Cell Rep.* **31**, 107808 (2020).
- Deshmukh, A. S. et al. Deep muscle-proteomic analysis of freeze-dried human muscle biopsies reveals fiber type-specific adaptations to exercise training. *Nat. Commun.* **12**, 304 (2021).
This study highlights notable shared and distinct adaptations in the proteomes of type I and type II fibres after a period of endurance training.
- Granata, C. et al. High-intensity training induces non-stoichiometric changes in the mitochondrial proteome of human skeletal muscle without reorganisation of respiratory chain content. *Nat. Commun.* **12**, 7056 (2021).
This study interrogates the muscle mitochondrial proteome at multiple time points during a periodized high-intensity training intervention.
- Laukkanen, J. A. et al. Long-term change in cardiorespiratory fitness and all-cause mortality: a population-based follow-up study. *Mayo Clin. Proc.* **91**, 1183–1188 (2016).
- Garcia-Hermoso, A. et al. Muscular strength as a predictor of all-cause mortality in an apparently healthy population: a systematic review and meta-analysis of data from approximately 2 million men and women. *Arch. Phys. Med. Rehabil.* **99**, 2100–2113.e5 (2018).
- Coleman, C. J., McDonough, D. J., Pope, Z. C. & Pope, C. A. Dose-response association of aerobic and muscle-strengthening physical activity with mortality: a national cohort study of 416 420 US adults. *Br. J. Sports Med.* <https://doi.org/10.1136/bjsports-2022-105519> (2022).
- Contrepour, K. et al. Molecular choreography of acute exercise. *Cell* **181**, 1112–1130.e16 (2020).
This study is a multi-omics analysis of human plasma and peripheral blood mononuclear cells at multiple time points across a 1-h recovery period immediately following a standardized exercise bout.
- Sato, S. et al. Atlas of exercise metabolism reveals time-dependent signatures of metabolic homeostasis. *Cell Metab.* **34**, 329–345.e8 (2022).
In this study, mass spectrometry-based metabolomics is used to characterize time-of-day differences in metabolic programming in several mouse tissues in response to an acute bout of exercise, including the net uptake and release of metabolites in hindlimb muscles and liver.
- Murgia, M. et al. Protein profile of fiber types in human skeletal muscle: a single-fiber proteomics study. *Skelet. Muscle* **11**, 24 (2021).
This study interrogates the proteome of single muscle fibres from young, healthy individuals and provides a useful resource for basal proteomic comparisons between type I and type II fibres.
- Bloemberg, D. & Quadrilatero, J. Rapid determination of myosin heavy chain expression in rat, mouse, and human skeletal muscle using multicolor immunofluorescence analysis. *PLoS ONE* **7**, e35273 (2012).
This paper provides a valuable resource for myosin fibre type and enzymatic profile comparisons, and identifies notable differences between muscles and species.
- Simoneau, J. A. & Bouchard, C. Human variation in skeletal muscle fiber-type proportion and enzyme activities. *Am. J. Physiol.* **257**, E567–E572 (1989).
- Pellegrino, M. A. et al. Orthologous myosin isoforms and scaling of shortening velocity with body size in mouse, rat, rabbit and human muscles. *J. Physiol.* **546**, 677–689 (2003).
- Bottinelli, R., Pellegrino, M. A., Canepari, M., Rossi, R. & Reggiani, C. Specific contributions of various muscle fibre types to human muscle performance: an in vitro study. *J. Electromyogr. Kinesiol.* **9**, 87–95 (1999).
- Dos Santos, M. et al. Single-nucleus RNA-seq and FISH identify coordinated transcriptional activity in mammalian myofibers. *Nat. Commun.* **11**, 5102 (2020).
This is one of the first studies to perform single myonuclear RNA-sequencing, revealing heterogeneity between myonuclei within a given muscle fibre.
- Dos Santos, M. et al. A fast myosin super enhancer dictates muscle fiber phenotype through competitive interactions with myosin genes. *Nat. Commun.* **13**, 1039 (2022).
This study uses a 'rainbow' transgenic mouse model of the fast-type myosin locus to determine that competitive promoter–super enhancer interactions govern fast-twitch myosin isoform expression in skeletal muscle.
- Roman, W. et al. Myofibril contraction and crosslinking drive nuclear movement to the periphery of skeletal muscle. *Nat. Cell Biol.* **19**, 1189–1201 (2017).
- Wen, Y. et al. Myonuclear transcriptional dynamics in response to exercise following satellite cell depletion. *iScience* **24**, 102838 (2021).
- Kim, M. et al. Single-nucleus transcriptomics reveals functional compartmentalization in syncytial skeletal muscle cells. *Nat. Commun.* **11**, 6375 (2020).
- D'Ercole, C. et al. Spatially resolved transcriptomics reveals innervation-responsive functional clusters in skeletal muscle. *Cell Rep.* **41**, 111861 (2022).
- Batthey, E. et al. Myonuclear alterations associated with exercise are independent of age in humans. *J. Physiol.* <https://doi.org/10.1113/JP284128> (2023).

33. Viggers, M. R. et al. Adaptation of the transcriptional response to resistance exercise over 4 weeks of daily training. *FASEB J.* **37**, e22686 (2023).
34. Lexell, J., Taylor, C. C. & Sjöström, M. What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. *J. Neurol. Sci.* **84**, 275–294 (1988).
This classic study establishes fibre type differences in human skeletal muscle during healthy ageing, thereby providing insights on the aetiology of sarcopenia.
35. Nielsen, J., Holmberg, H. C., Schroder, H. D., Saltin, B. & Ortenblad, N. Human skeletal muscle glycogen utilization in exhaustive exercise: role of subcellular localization and fibre type. *J. Physiol.* **589**, 2871–2885 (2011).
36. Murgija, M. et al. Single muscle fiber proteomics reveals fiber-type-specific features of human muscle aging. *Cell Rep.* **19**, 2396–2409 (2017).
37. Luo, L. et al. Spatial metabolomics reveals skeletal myofiber subtypes. *Sci. Adv.* **9**, eadd0455 (2023).
38. Medler, S. Mixing it up: the biological significance of hybrid skeletal muscle fibers. *J. Exp. Biol.* **222**, jeb200832 (2019).
39. Horwath, O. et al. Variability in vastus lateralis fiber type distribution, fiber size, and myonuclear content along and between the legs. *J. Appl. Physiol.* **131**, 158–173 (2021).
40. Murach, K. A. et al. Fiber typing human skeletal muscle with fluorescent immunohistochemistry. *J. Appl. Physiol.* **127**, 1632–1639 (2019).
41. Bathgate, K. E. et al. Muscle health and performance in monozygotic twins with 30 years of discordant exercise habits. *Eur. J. Appl. Physiol.* **118**, 2097–2110 (2018).
This study of monozygotic twins suggests that fibre type plasticity in response to lifelong endurance training is greater than previously appreciated.
42. Malisoux, L., Francaux, M., Nielsen, H. & Theisen, D. Stretch-shortening cycle exercises: an effective training paradigm to enhance power output of human single muscle fibers. *J. Appl. Physiol.* **100**, 771–779 (2006).
43. Sakakibara, I., Santolini, M., Ferry, A., Hakim, V. & Maire, P. Six homeoproteins and a linc-RNA at the fast MYH locus lock fast myofiber terminal phenotype. *PLoS Genet.* **10**, e1004386 (2014).
44. Trappe, S. et al. Single muscle fiber adaptations with marathon training. *J. Appl. Physiol.* **101**, 721–727 (2006).
45. Williamson, D. L., Gallagher, P. M., Carroll, C. C., Raue, U. & Trappe, S. W. Reduction in hybrid single muscle fiber proportions with resistance training in humans. *J. Appl. Physiol.* **91**, 1955–1961 (2001).
46. Long, K. et al. Identification of enhancers responsible for the coordinated expression of myosin heavy chain isoforms in skeletal muscle. *BMC Genomics* **23**, 519 (2022).
This study, published shortly after Dos Santos et al. (2022), also identifies the fast-myosin super enhancer, as well as enhancers of slow-type myosin.
47. Balagopal, P., Schimke, J. C., Ades, P., Adey, D. & Nair, K. S. Age effect on transcript levels and synthesis rate of muscle MHC and response to resistance exercise. *Am. J. Physiol. Endocrinol. Metab.* **280**, E203–E208 (2001).
48. Lin, J. et al. Transcriptional co-activator PGC-1 α drives the formation of slow-twitch muscle fibres. *Nature* **418**, 797–801 (2002).
49. Kuhnen, G. et al. Genes whose gain or loss of function changes type 1, 2A, 2X, or 2B muscle fibre proportions in mice — a systematic review. *Int. J. Mol. Sci.* **23**, 12933 (2022).
50. Street, S. F. Lateral transmission of tension in frog myofibers: a myofibrillar network and transverse cytoskeletal connections are possible transmitters. *J. Cell. Physiol.* **114**, 346–364 (1983).
51. Willingham, T. B., Kim, Y., Lindberg, E., Bleck, C. K. E. & Glancy, B. The unified myofibrillar matrix for force generation in muscle. *Nat. Commun.* **11**, 3722 (2020).
The microscopic analysis of muscle myofibrils in this study provides a new paradigm for the organization of muscle contractile apparatus.
52. Ajayi, P. T. et al. Regulation of the evolutionarily conserved muscle myofibrillar matrix by cell type dependent and independent mechanisms. *Nat. Commun.* **13**, 2661 (2022).
53. Katti, P. et al. Mitochondrial network configuration influences sarcomere and myosin filament structure in striated muscles. *Nat. Commun.* **13**, 6058 (2022).
54. Krustup, P., Ferguson, R. A., Kjaer, M. & Bangsbo, J. ATP and heat production in human skeletal muscle during dynamic exercise: higher efficiency of anaerobic than aerobic ATP resynthesis. *J. Physiol.* **549**, 255–269 (2003).
55. Zagatto, A. M. et al. Impacts of high-intensity exercise on the metabolomics profile of human skeletal muscle tissue. *Scand. J. Med. Sci. Sports* **32**, 402–413 (2022).
This study is an analysis of the immediate post-exercise skeletal muscle metabolome in response to an exhaustive bout of high-intensity cycling.
56. Bleck, C. K. E., Kim, Y., Willingham, T. B. & Glancy, B. Subcellular connectomic analyses of energy networks in striated muscle. *Nat. Commun.* **9**, 5111 (2018).
57. Glancy, B. et al. Mitochondrial reticulum for cellular energy distribution in muscle. *Nature* **523**, 617–620 (2015).
This work advances the understanding of how membrane potential is dispersed across the mitochondrial reticulum to facilitate efficient ATP distribution in muscle.
58. Glancy, B. et al. Power grid protection of the muscle mitochondrial reticulum. *Cell Rep.* **19**, 487–496 (2017).
59. Vincent, A. E. et al. Quantitative 3D mapping of the human skeletal muscle mitochondrial network. *Cell Rep.* **26**, 996–1009.e4 (2019).
60. Caffrey, B. J. et al. Semi-automated 3D segmentation of human skeletal muscle using focused ion beam-scanning electron microscopic images. *J. Struct. Biol.* **207**, 1–11 (2019).
61. Greggio, C. et al. Enhanced respiratory chain supercomplex formation in response to exercise in human skeletal muscle. *Cell Metab.* **25**, 301–311 (2017).
62. Cogliati, S. et al. Mitochondrial cristae shape determines respiratory chain supercomplexes assembly and respiratory efficiency. *Cell* **155**, 160–171 (2013).
63. Balsa, E. et al. ER and nutrient stress promote assembly of respiratory chain supercomplexes through the PERK-eIF2 α axis. *Mol. Cell* **74**, 877–890.e6 (2019).
64. Nielsen, J. et al. Plasticity in mitochondrial cristae density allows metabolic capacity modulation in human skeletal muscle. *J. Physiol.* **595**, 2839–2847 (2017).
65. Hentila, J. et al. Autophagy is induced by resistance exercise in young men, but unfolded protein response is induced regardless of age. *Acta Physiol.* **224**, e13069 (2018).
66. Heymsfield, S. B. et al. Human energy expenditure: advances in organ-tissue prediction models. *Obes. Rev.* **19**, 1177–1188 (2018).
67. Phung, L. A., Foster, A. D., Miller, M. S., Lowe, D. A. & Thomas, D. D. Super-relaxed state of myosin in human skeletal muscle is fiber-type dependent. *Am. J. Physiol. Cell Physiol.* **319**, C1158–C1162 (2020).
68. Linari, M. et al. Force generation by skeletal muscle is controlled by mechanosensing in myosin filaments. *Nature* **528**, 276–279 (2015).
69. Gaitanos, G. C., Williams, C., Boobis, L. H. & Brooks, S. Human muscle metabolism during intermittent maximal exercise. *J. Appl. Physiol.* **75**, 712–719 (1993).
70. Parolin, M. L. et al. Regulation of skeletal muscle glycogen phosphorylase and PDH during maximal intermittent exercise. *Am. J. Physiol.* **277**, E890–E900 (1999).
71. Romijn, J. A. et al. Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am. J. Physiol.* **265**, E380–E391 (1993).
72. van Loon, L. J., Greenhaff, P. L., Constantini-Teodosiu, D., Saris, W. H. & Wagenmakers, A. J. The effects of increasing exercise intensity on muscle fuel utilisation in humans. *J. Physiol.* **536**, 295–304 (2001).
73. Duchateau, J. & Enoka, R. M. Human motor unit recordings: origins and insight into the integrated motor system. *Brain Res.* **1409**, 42–61 (2011).
74. Stienen, G. J., Kiers, J. L., Bottinelli, R. & Reggiani, C. Myofibrillar ATPase activity in skinned human skeletal muscle fibres: fibre type and temperature dependence. *J. Physiol.* **493**, 299–307 (1996).
75. Greenhaff, P. L. et al. The metabolic responses of human type I and II muscle fibres during maximal treadmill sprinting. *J. Physiol.* **478**, 149–155 (1994).
76. Koh, H. E., Nielsen, J., Saltin, B., Holmberg, H. C. & Ortenblad, N. Pronounced limb and fibre type differences in subcellular lipid droplet content and distribution in elite skiers before and after exhaustive exercise. *J. Physiol.* **595**, 5781–5795 (2017).
77. de Almeida, M. E. et al. Altered intramuscular network of lipid droplets and mitochondria in type 2 diabetes. *Am. J. Physiol. Cell Physiol.* **324**, C39–C57 (2023).
78. Daemen, S. et al. Distinct lipid droplet characteristics and distribution unmask the apparent contradiction of the athlete's paradox. *Mol. Metab.* **17**, 71–81 (2018).
79. Brooks, G. A. & Mercier, J. Balance of carbohydrate and lipid utilization during exercise: the "crossover" concept. *J. Appl. Physiol.* **76**, 2253–2261 (1994).
80. Wust, R. C. et al. Kinetic control of oxygen consumption during contractions in self-perfused skeletal muscle. *J. Physiol.* **589**, 3995–4009 (2011).
81. Ferguson, B. S. et al. Lactate metabolism: historical context, prior misinterpretations, and current understanding. *Eur. J. Appl. Physiol.* **118**, 691–728 (2018).
82. Wescott, A. P., Kao, J. P. Y., Lederer, W. J. & Boyman, L. Voltage-energized calcium-sensitive ATP production by mitochondria. *Nat. Metab.* **1**, 975–984 (2019).
83. Seiler, S. E. et al. Carnitine acetyltransferase mitigates metabolic inertia and muscle fatigue during exercise. *Cell Metab.* **22**, 65–76 (2015).
84. Mancilla, R. F. et al. Skeletal muscle mitochondrial inertia is associated with carnitine acetyltransferase activity and physical function in humans. *JCI Insight* **8**, e163855 (2023).
85. Timmons, J. A. et al. Substrate availability limits human skeletal muscle oxidative ATP regeneration at the onset of ischemic exercise. *J. Clin. Invest.* **101**, 79–85 (1998).
86. Walsh, B. et al. The role of phosphorylcreatine and creatine in the regulation of mitochondrial respiration in human skeletal muscle. *J. Physiol.* **537**, 971–978 (2001).
87. Roman, B. B., Meyer, R. A. & Wiseman, R. W. Phosphocreatine kinetics at the onset of contractions in skeletal muscle of MM creatine kinase knockout mice. *Am. J. Physiol. Cell Physiol.* **283**, C1776–C1783 (2002).
88. Katz, A., Broberg, S., Sahlin, K. & Wahren, J. Leg glucose uptake during maximal dynamic exercise in humans. *Am. J. Physiol.* **251**, E65–E70 (1986).
89. Katz, A. A century of exercise physiology: key concepts in regulation of glycogen metabolism in skeletal muscle. *Eur. J. Appl. Physiol.* **122**, 1751–1772 (2022).
90. Blazev, R. et al. Phosphoproteomics of three exercise modalities identifies canonical signaling and C18ORF25 as an AMPK substrate regulating skeletal muscle function. *Cell Metab.* **34**, 1561–1577.e1569 (2022).
Building on work from Hoffman et al. (2015), this study compares the phosphoproteomic impact of an acute bout of endurance, resistance or sprinting exercise across a 3-h post-exercise recovery period.
91. Watt, M. J. et al. Regulation of HSL serine phosphorylation in skeletal muscle and adipose tissue. *Am. J. Physiol. Endocrinol. Metab.* **290**, E500–E508 (2006).
92. Richter, E. A., Ruderman, N. B., Gavras, H., Belur, E. R. & Galbo, H. Muscle glycogenolysis during exercise: dual control by epinephrine and contractions. *Am. J. Physiol.* **242**, E25–E32 (1982).
93. Hureau, T. J. et al. On the role of skeletal muscle acidosis and inorganic phosphates as determinants of central and peripheral fatigue: a ^{31}P -MRS study. *J. Physiol.* **600**, 3069–3081 (2022).
94. Cheng, A. J., Place, N. & Westerblad, H. Molecular basis for exercise-induced fatigue: the importance of strictly controlled cellular Ca(2+) handling. *Cold Spring Harb. Perspect. Med.* **8**, a029710 (2018).

95. Rogatzki, M. J., Ferguson, B. S., Goodwin, M. L. & Gladden, L. B. Lactate is always the end product of glycolysis. *Front. Neurosci.* **9**, 22 (2015).
96. Hokken, R. et al. Subcellular localization- and fibre type-dependent utilization of muscle glycogen during heavy resistance exercise in elite power and Olympic weightlifters. *Acta Physiol.* **231**, e13561 (2021).
97. Vigh-Larsen, J. F. et al. Fibre type- and localisation-specific muscle glycogen utilisation during repeated high-intensity intermittent exercise. *J. Physiol.* **600**, 4713–4730 (2022).
98. Sidhu, S. K. et al. Fatigue-related group III/IV muscle afferent feedback facilitates intracortical inhibition during locomotor exercise. *J. Physiol.* **596**, 4789–4801 (2018).
99. Ducrocq, G. P. & Blain, G. M. Relationship between neuromuscular fatigue, muscle activation and the work done above the critical power during severe-intensity exercise. *Exp. Physiol.* **107**, 312–325 (2022).
100. Goncalves, R. L., Quinlan, C. L., Perevoshchikova, I. V., Hey-Mogensen, M. & Brand, M. D. Sites of superoxide and hydrogen peroxide production by muscle mitochondria assessed ex vivo under conditions mimicking rest and exercise. *J. Biol. Chem.* **290**, 209–227 (2015).
101. Henriquez-Olguin, C. et al. Cytosolic ROS production by NADPH oxidase 2 regulates muscle glucose uptake during exercise. *Nat. Commun.* **10**, 4623 (2019).
102. Sakellariou, G. K. et al. Studies of mitochondrial and nonmitochondrial sources implicate nicotinamide adenine dinucleotide phosphate oxidase(s) in the increased skeletal muscle superoxide generation that occurs during contractile activity. *Antioxid. Redox Signal.* **18**, 603–621 (2013).
103. Sun, Q. A. et al. Oxygen-coupled redox regulation of the skeletal muscle ryanodine receptor-Ca²⁺ release channel by NADPH oxidase 4. *Proc. Natl Acad. Sci. USA* **108**, 16098–16103 (2011).
104. Xirouchaki, C. E. et al. Skeletal muscle NOX4 is required for adaptive responses that prevent insulin resistance. *Sci. Adv.* **7**, eabl4988 (2021).
105. Bouviere, J. et al. Exercise-stimulated ROS sensitive signaling pathways in skeletal muscle. *Antioxidants* **10**, 537 (2021).
106. Ristow, M. et al. Antioxidants prevent health-promoting effects of physical exercise in humans. *Proc. Natl Acad. Sci. USA* **106**, 8665–8670 (2009).
107. Clifford, T., Jeffries, O., Stevenson, E. J. & Davies, K. A. B. The effects of vitamin C and E on exercise-induced physiological adaptations: a systematic review and Meta-analysis of randomized controlled trials. *Crit. Rev. Food Sci. Nutr.* **60**, 3669–3679 (2020).
108. Gonzalez, J. T. et al. Ingestion of glucose or sucrose prevents liver but not muscle glycogen depletion during prolonged endurance-type exercise in trained cyclists. *Am. J. Physiol. Endocrinol. Metab.* **309**, E1032–E1039 (2015).
109. Vincent, M. A. et al. Mixed meal and light exercise each recruit muscle capillaries in healthy humans. *Am. J. Physiol. Endocrinol. Metab.* **290**, E1191–E1197 (2006).
110. MacLean, D. A., Bangsbo, J. & Saltin, B. Muscle interstitial glucose and lactate levels during dynamic exercise in humans determined by microdialysis. *J. Appl. Physiol.* **87**, 1483–1490 (1999).
111. Chow, L. S. et al. Exerkines in health, resilience and disease. *Nat. Rev. Endocrinol.* **18**, 273–289 (2022).
112. Ryder, J. W. et al. Postexercise glucose uptake and glycogen synthesis in skeletal muscle from GLUT4-deficient mice. *FASEB J.* **13**, 2246–2256 (1999).
113. Sylow, L., Kleinert, M., Richter, E. A. & Jensen, T. E. Exercise-stimulated glucose uptake – regulation and implications for glycaemic control. *Nat. Rev. Endocrinol.* **13**, 133–148 (2017).
114. Sylow, L. et al. Rac1 and AMPK account for the majority of muscle glucose uptake stimulated by ex vivo contraction but not in vivo exercise. *Diabetes* **66**, 1548–1559 (2017).
115. Kjobsted, R. et al. AMPK and TBC1D1 regulate muscle glucose uptake after, but not during, exercise and contraction. *Diabetes* **68**, 1427–1440 (2019).
116. Kjobsted, R. et al. Enhanced muscle insulin sensitivity after contraction/exercise is mediated by AMPK. *Diabetes* **66**, 598–612 (2017).
117. Fritzen, A. M. et al. 5'-AMP activated protein kinase α_2 controls substrate metabolism during post-exercise recovery via regulation of pyruvate dehydrogenase kinase 4. *J. Physiol.* **593**, 4765–4780 (2015).
118. Ouyang, Q. et al. Rab8a as a mitochondrial receptor for lipid droplets in skeletal muscle. *Dev. Cell* **58**, 289–305.e6 (2023).
119. Glancy, B. et al. Mitochondrial lactate metabolism: history and implications for exercise and disease. *J. Physiol.* **599**, 863–888 (2021).
120. Rothschild, J. A., Kilding, A. E., Stewart, T. & Plews, D. J. Factors influencing substrate oxidation during submaximal cycling: a modelling analysis. *Sports Med.* **52**, 2775–2795 (2022).
121. King, A., Helms, E., Zinn, C. & Jukic, I. The ergogenic effects of acute carbohydrate feeding on resistance exercise performance: a systematic review and meta-analysis. *Sports Med.* **52**, 2691–2712 (2022).
122. Hartley, C., Carr, A., Bowe, S. J., Bredie, W. L. P. & Keast, R. S. J. Maltodextrin-based carbohydrate oral rinsing and exercise performance: systematic review and meta-analysis. *Sports Med.* **52**, 1833–1862 (2022).
123. Gant, N., Stinear, C. M. & Byblow, W. D. Carbohydrate in the mouth immediately facilitates motor output. *Brain Res.* **1350**, 151–158 (2010).
124. Morville, T., Sahl, R. E., Moritz, T., Helge, J. W. & Clemmensen, C. Plasma metabolome profiling of resistance exercise and endurance exercise in humans. *Cell Rep.* **33**, 108554 (2020).
125. Alsted, T. J. et al. Contraction-induced lipolysis is not impaired by inhibition of hormone-sensitive lipase in skeletal muscle. *J. Physiol.* **591**, 5141–5155 (2013).
126. Dube, J. J. et al. Adipose triglyceride lipase deletion from adipocytes, but not skeletal myocytes, impairs acute exercise performance in mice. *Am. J. Physiol. Endocrinol. Metab.* **308**, E879–E890 (2015).
127. Watt, M. J., Heigenhauser, G. J. & Spriet, L. L. Effects of dynamic exercise intensity on the activation of hormone-sensitive lipase in human skeletal muscle. *J. Physiol.* **547**, 301–308 (2003).
128. Donsmark, M., Langfort, J., Holm, C., Ploug, T. & Galbo, H. Contractions induce phosphorylation of the AMPK site Ser565 in hormone-sensitive lipase in muscle. *Biochem. Biophys. Res. Commun.* **316**, 867–871 (2004).
129. Prats, C. et al. Decrease in intramuscular lipid droplets and translocation of HSL in response to muscle contraction and epinephrine. *J. Lipid Res.* **47**, 2392–2399 (2006).
130. Stokic, J. R., Abbott, G., Howlett, K. F., Hamilton, D. L. & Shaw, C. S. Intramuscular lipid utilization during exercise: a systematic review, meta-analysis, and meta-regression. *J. Appl. Physiol.* **134**, 581–592 (2023).
131. Ghafouri, K. et al. Moderate exercise increases affinity of large very low-density lipoproteins for hydrolysis by lipoprotein lipase. *J. Clin. Endocrinol. Metab.* **100**, 2205–2213 (2015).
132. Jain, S. S. et al. Additive effects of insulin and muscle contraction on fatty acid transport and fatty acid transporters, FAT/CD36, FABPm, FATP1, 4 and 6. *FEBS Lett.* **583**, 2294–2300 (2009).
133. Nickerson, J. G. et al. Greater transport efficiencies of the membrane fatty acid transporters FAT/CD36 and FATP4 compared with FABPm and FATP1 and differential effects on fatty acid esterification and oxidation in rat skeletal muscle. *J. Biol. Chem.* **284**, 16522–16530 (2009).
134. Jeppesen, J. et al. Contraction-induced skeletal muscle FAT/CD36 trafficking and FA uptake is AMPK independent. *J. Lipid Res.* **52**, 699–711 (2011).
135. Abbott, M. J., Edelman, A. M. & Turcotte, L. P. CaMKK is an upstream signal of AMP-activated protein kinase in regulation of substrate metabolism in contracting skeletal muscle. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **297**, R1724–R1732 (2009).
136. Turcotte, L. P., Raney, M. A. & Todd, M. K. ERK1/2 inhibition prevents contraction-induced increase in plasma membrane FAT/CD36 content and FA uptake in rodent muscle. *Acta Physiol. Scand.* **184**, 131–139 (2005).
137. Holloway, G. P. et al. Fatty acid binding protein facilitates sarcolemmal fatty acid transport but not mitochondrial oxidation in rat and human skeletal muscle. *J. Physiol.* **582**, 393–405 (2007).
138. Sacchetti, M., Saltin, B., Osada, T. & van Hall, G. Intramuscular fatty acid metabolism in contracting and non-contracting human skeletal muscle. *J. Physiol.* **540**, 387–395 (2002).
139. Hall, A. M., Wiczler, B. M., Herrmann, T., Stremmel, W. & Bernlohr, D. A. Enzymatic properties of purified murine fatty acid transport protein 4 and analysis of acyl-CoA synthetase activities in tissues from FATP4 null mice. *J. Biol. Chem.* **280**, 11948–11954 (2005).
140. Li, L. O. et al. Compartmentalized acyl-CoA metabolism in skeletal muscle regulates systemic glucose homeostasis. *Diabetes* **64**, 23–35 (2015).
141. Chen, S., Zhou, L., Sun, J., Qu, Y. & Chen, M. The role of cAMP-PKA pathway in lactate-induced intramuscular triglyceride accumulation and mitochondria content increase in mice. *Front. Physiol.* **12**, 709135 (2021).
142. Constantin-Teodosiu, D., Howell, S. & Greenhaff, P. L. Carnitine metabolism in human muscle fiber types during submaximal dynamic exercise. *J. Appl. Physiol.* **80**, 1061–1064 (1996).
143. Hostrup, M. et al. High-intensity interval training remodels the proteome and acetylome of human skeletal muscle. *eLife* **11**, e69802 (2022).
144. Adamovich, Y. et al. Clock proteins and training modify exercise capacity in a daytime-dependent manner. *Proc. Natl Acad. Sci. USA* **118**, e210115118 (2021).
145. Bano-Otalora, B. et al. Bright daytime light enhances circadian amplitude in a diurnal mammal. *Proc. Natl Acad. Sci. USA* **118**, e2100094118 (2021).
146. Martin, R. A., Viggers, M. R. & Esser, K. A. Metabolism and exercise: the skeletal muscle clock takes centre stage. *Nat. Rev. Endocrinol.* <https://doi.org/10.1038/s41574-023-00805-8> (2023).
147. Dyar, K. A. et al. Muscle insulin sensitivity and glucose metabolism are controlled by the intrinsic muscle clock. *Mol. Metab.* **3**, 29–41 (2014).
148. Dyar, K. A. et al. Transcriptional programming of lipid and amino acid metabolism by the skeletal muscle circadian clock. *PLoS Biol.* **16**, e2005886 (2018).
149. Hodge, B. A. et al. The endogenous molecular clock orchestrates the temporal separation of substrate metabolism in skeletal muscle. *Skelet. Muscle* **5**, 17 (2015).
150. Harfmann, B. D. et al. Muscle-specific loss of Bmal1 leads to disrupted tissue glucose metabolism and systemic glucose homeostasis. *Skelet. Muscle* **6**, 12 (2016).
151. Thomas, J. M. et al. Circadian rhythm phase shifts caused by timed exercise vary with chronotype. *JCI Insight* **5**, e134270 (2020).
152. Sun, S. et al. A single-cell transcriptomic atlas of exercise-induced anti-inflammatory and geroprotective effects across the body. *Innovation* **4**, 100380 (2023).
153. Small, L. et al. Contraction influences Per2 gene expression in skeletal muscle through a calcium-dependent pathway. *J. Physiol.* **598**, 5739–5752 (2020).
154. Zambon, A. C. et al. Time- and exercise-dependent gene regulation in human skeletal muscle. *Genome Biol.* **4**, R61 (2003).
155. Dyar, K. A. et al. The calcineurin-NFAT pathway controls activity-dependent circadian gene expression in slow skeletal muscle. *Mol. Metab.* **4**, 823–833 (2015).
156. Gabriel, B. M. & Zierath, J. R. Circadian rhythms and exercise – re-setting the clock in metabolic disease. *Nat. Rev. Endocrinol.* **15**, 197–206 (2019).
157. Harmsen, J. F. et al. Circadian misalignment disturbs the skeletal muscle lipidome in healthy young men. *FASEB J.* **35**, e21611 (2021).

158. Wefers, J. et al. Circadian misalignment induces fatty acid metabolism gene profiles and compromises insulin sensitivity in human skeletal muscle. *Proc. Natl Acad. Sci. USA* **115**, 7789–7794 (2018).
159. Morris, C. J., Purvis, T. E., Hu, K. & Scheer, F. A. Circadian misalignment increases cardiovascular disease risk factors in humans. *Proc. Natl Acad. Sci. USA* **113**, E1402–E1411 (2016).
160. Saner, N. J. et al. Exercise mitigates sleep-loss-induced changes in glucose tolerance, mitochondrial function, sarcoplasmic protein synthesis, and diurnal rhythms. *Mol. Metab.* **43**, 101110 (2021).
161. Sato, S. et al. Time of exercise specifies the impact on muscle metabolic pathways and systemic energy homeostasis. *Cell Metab.* **30**, 92–110.e4 (2019).
162. Mancilla, R. et al. Exercise training elicits superior metabolic effects when performed in the afternoon compared to morning in metabolically compromised humans. *Physiol. Rep.* **8**, e14669 (2021).
163. van der Velde, J. et al. Timing of physical activity in relation to liver fat content and insulin resistance. *Diabetologia* **66**, 461–471 (2023).
164. Savikj, M. et al. Afternoon exercise is more efficacious than morning exercise at improving blood glucose levels in individuals with type 2 diabetes: a randomised crossover trial. *Diabetologia* **62**, 233–237 (2019).
165. Feng, H. et al. Associations of timing of physical activity with all-cause and cause-specific mortality in a prospective cohort study. *Nat. Commun.* **14**, 930 (2023).
166. Fischer, D., Lombardi, D. A., Marucci-Wellman, H. & Roenneberg, T. Chronotypes in the US — influence of age and sex. *PLoS ONE* **12**, e0178782 (2017).
167. Amar, D. et al. Time trajectories in the transcriptomic response to exercise — a meta-analysis. *Nat. Commun.* **12**, 3471 (2021).
168. Pillon, N. J. et al. Distinctive exercise-induced inflammatory response and exerkin induction in skeletal muscle of people with type 2 diabetes. *Sci. Adv.* **8**, eabo3192 (2022).
169. Makhnovskii, P. A. et al. Alternative transcription start sites contribute to acute-stress-induced transcriptome response in human skeletal muscle. *Hum. Genomics* **16**, 24 (2022).
170. Hoffman, N. J. et al. Global phosphoproteomic analysis of human skeletal muscle reveals a network of exercise-regulated kinases and AMPK substrates. *Cell Metab.* **22**, 922–935 (2015).
- This seminal study investigates the exercise-induced human muscle phosphoproteome.**
171. Nelson, M. E. et al. Phosphoproteomics reveals conserved exercise-stimulated signaling and AMPK regulation of store-operated calcium entry. *EMBO J.* **38**, e102578 (2019).
172. You, J. S. et al. The role of raptor in the mechanical load-induced regulation of mTOR signaling, protein synthesis, and skeletal muscle hypertrophy. *FASEB J.* **33**, 4021–4034 (2019).
173. You, J. S. et al. A DGK ζ -FoxO-ubiquitin proteolytic axis controls fiber size during skeletal muscle remodeling. *Sci. Signal.* **11**, eaa06847 (2018).
174. Song, Z. et al. Resistance exercise initiates mechanistic target of rapamycin (mTOR) translocation and protein complex co-localisation in human skeletal muscle. *Sci. Rep.* **7**, 5028 (2017).
175. Hodson, N., Mazzulla, M., Holowaty, M. N. H., Kumbhare, D. & Moore, D. R. RPS6 phosphorylation occurs to a greater extent in the periphery of human skeletal muscle fibers, near focal adhesions, after anabolic stimuli. *Am. J. Physiol. Cell Physiol.* **322**, C94–C110 (2022).
176. Abou Sawan, S. et al. Trained integrated postexercise myofibrillar protein synthesis rates correlate with hypertrophy in young males and females. *Med. Sci. Sports Exerc.* **54**, 953–964 (2022).
177. Needham, E. J. et al. Personalized phosphoproteomics identifies functional signaling. *Nat. Biotechnol.* **40**, 576–584 (2022).
178. Steinert, N. D. et al. Mapping of the contraction-induced phosphoproteome identifies TRIM28 as a significant regulator of skeletal muscle size and function. *Cell Rep.* **34**, 108796 (2021).
179. Nordgaard, C. et al. ZAK β is activated by cellular compression and mediates contraction-induced MAP kinase signaling in skeletal muscle. *EMBO J.* **41**, e111650 (2022).
180. West, D. W. et al. Acute resistance exercise activates rapamycin-sensitive and -insensitive mechanisms that control translational activity and capacity in skeletal muscle. *J. Physiol.* **594**, 453–468 (2016).
181. Lessard, S. J. et al. JNK regulates muscle remodeling via myostatin/SMAD inhibition. *Nat. Commun.* **9**, 3030 (2018).
182. MacKenzie, M. G., Hamilton, D. L., Pepin, M., Patton, A. & Baar, K. Inhibition of myostatin signaling through Notch activation following acute resistance exercise. *PLoS ONE* **8**, e68743 (2013).
183. Pillon, N. J. et al. Transcriptomic profiling of skeletal muscle adaptations to exercise and inactivity. *Nat. Commun.* **11**, 470 (2020).
- This meta-analysis curates an extensive (and frequently updated) library of the skeletal muscle transcriptomic response to exercise across human demographics and exercise modalities.**
184. Schumann, M. et al. Compatibility of concurrent aerobic and strength training for skeletal muscle size and function: an updated systematic review and meta-analysis. *Sports Med.* **52**, 601–612 (2022).
185. Lundberg, T. R., Feuerbacher, J. F., Sunkeler, M. & Schumann, M. The effects of concurrent aerobic and strength training on muscle fiber hypertrophy: a systematic review and meta-analysis. *Sports Med.* **52**, 2391–2403 (2022).
186. Kaiser, M. S. et al. Dual roles of mTORC1-dependent activation of the ubiquitin-proteasome system in muscle proteostasis. *Commun. Biol.* **5**, 1141 (2022).
187. Drake, J. C. et al. Mitochondria-localized AMPK responds to local energetics and contributes to exercise and energetic stress-induced mitophagy. *Proc. Natl Acad. Sci. USA* **118**, e2025932118 (2021).
188. Laker, R. C. et al. AMPK phosphorylation of Ulk1 is required for targeting of mitochondria to lysosomes in exercise-induced mitophagy. *Nat. Commun.* **8**, 548 (2017).
189. Toyama, E. Q. et al. Metabolism. AMP-activated protein kinase mediates mitochondrial fission in response to energy stress. *Science* **351**, 275–281 (2016).
190. VerPlank, J. J. S., Lokireddy, S., Zhao, J. & Goldberg, A. L. 26S Proteasomes are rapidly activated by diverse hormones and physiological states that raise cAMP and cause Rpn6 phosphorylation. *Proc. Natl Acad. Sci. USA* **116**, 4228–4237 (2019).
191. Parker, B. L., Kiens, B., Wojtaszewski, J. F. P., Richter, E. A. & James, D. E. Quantification of exercise-regulated ubiquitin signaling in human skeletal muscle identifies protein modification cross talk via NEDDylation. *FASEB J.* **34**, 5906–5916 (2020).
192. Seaborne, R. A. & Sharples, A. P. The interplay between exercise metabolism, epigenetics, and skeletal muscle remodeling. *Exerc. Sport Sci. Rev.* **48**, 188–200 (2020).
193. McGee, S. L., Fairlie, E., Garnham, A. P. & Hargreaves, M. Exercise-induced histone modifications in human skeletal muscle. *J. Physiol.* **587**, 5951–5958 (2009).
194. Backs, J., Song, K., Bezprozvannaya, S., Chang, S. & Olson, E. N. CaM kinase II selectively signals to histone deacetylase 4 during cardiomyocyte hypertrophy. *J. Clin. Invest.* **116**, 1853–1864 (2006).
195. Wu, X. et al. Local InsP3-dependent perinuclear Ca²⁺ signaling in cardiac myocyte excitation-transcription coupling. *J. Clin. Invest.* **116**, 675–682 (2006).
196. Solagna, F. et al. Exercise-dependent increases in protein synthesis are accompanied by chromatin modifications and increased MRTF-SRF signalling. *Acta Physiol.* **230**, e13496 (2020).
197. Yin, Y. et al. Impact of cytosine methylation on DNA binding specificities of human transcription factors. *Science* **356**, eaaj2239 (2017).
198. Figueiredo, V. C. et al. Genetic and epigenetic regulation of skeletal muscle ribosome biogenesis with exercise. *J. Physiol.* **599**, 3363–3384 (2021).
199. Seaborne, R. A. et al. Human skeletal muscle possesses an epigenetic memory of hypertrophy. *Sci. Rep.* **8**, 1898 (2018).
- This comprehensive interrogation of the resistance exercise-induced methylome provides an epigenetic hypothesis for ‘muscle memory’.**
200. Sexton, C. L. et al. Skeletal muscle DNA methylation and mRNA responses to a bout of higher versus lower load resistance exercise in previously trained men. *Cells* **12**, 263 (2023).
201. Lindholm, M. E. et al. An integrative analysis reveals coordinated reprogramming of the epigenome and the transcriptome in human skeletal muscle after training. *Epigenetics* **9**, 1557–1569 (2014).
202. Maasar, M. F. et al. The comparative methylome and transcriptome after change of direction compared to straight line running exercise in human skeletal muscle. *Front. Physiol.* **12**, 619447 (2021).
203. Galle, E. et al. H3K18 lactylation marks tissue-specific active enhancers. *Genome Biol.* **23**, 207 (2022).
204. Massart, J. et al. Endurance exercise training-responsive miR-19b-3p improves skeletal muscle glucose metabolism. *Nat. Commun.* **12**, 5948 (2021).
205. Bonilauro, B. & Dallagiovanna, B. Long non-coding RNAs are differentially expressed after different exercise training programs. *Front. Physiol.* **11**, 567614 (2020).
206. Vechetti, I. J. Jr. et al. Mechanical overload-induced muscle-derived extracellular vesicles promote adipose tissue lipolysis. *FASEB J.* **35**, e21644 (2021).
207. Xhuti, D., Nilsson, M. I., Manta, K., Tarnopolsky, M. A. & Nederveen, J. P. Circulating exosome-like vesicle and skeletal muscle microRNAs are altered with age and resistance training. *J. Physiol.* <https://doi.org/10.1113/jp282663> (2023).
208. Watanabe, S. et al. Skeletal muscle releases extracellular vesicles with distinct protein and microRNA signatures that function in the muscle microenvironment. *PNAS Nexus* **1**, pgac173 (2022).
209. Murach, K. A. et al. Fusion-independent satellite cell communication to muscle fibers during load-induced hypertrophy. *Function* **1**, zqaa009 (2020).
210. Fry, C. S., Kirby, T. J., Kosmac, K., McCarthy, J. J. & Peterson, C. A. Myogenic progenitor cells control extracellular matrix production by fibroblasts during skeletal muscle hypertrophy. *Cell Stem Cell* **20**, 56–69 (2017).
211. Albanese, M. et al. MicroRNAs are minor constituents of extracellular vesicles that are rarely delivered to target cells. *PLoS Genet.* **17**, e1009951 (2021).
212. Wohlwend, M. et al. The exercise-induced long noncoding RNA CYTOR promotes fast-twitch myogenesis in aging. *Sci. Transl. Med.* **13**, eabc7367 (2021).
213. Nelson, B. R. et al. A peptide encoded by a transcript annotated as long noncoding RNA enhances SERCA activity in muscle. *Science* **351**, 271–275 (2016).
214. Anderson, D. M. et al. A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. *Cell* **160**, 595–606 (2015).
215. Pinheiro, H. et al. mRNA distribution in skeletal muscle is associated with mRNA size. *J. Cell Sci.* **134**, jcs256388 (2021).
216. Denes, L. T., Kelley, C. P. & Wang, E. T. Microtubule-based transport is essential to distribute RNA and nascent protein in skeletal muscle. *Nat. Commun.* **12**, 6079 (2021).
- This study and Pinheiro et al. (2021) suggest that mRNAs are transported along microtubules in muscle, which has notable implications for the regulation of the ‘myonuclear domain’.**
217. Scarborough, E. A. et al. Microtubules orchestrate local translation to enable cardiac growth. *Nat. Commun.* **12**, 1547 (2021).
218. Taylor-Weiner, H. et al. Modeling the transport of nuclear proteins along single skeletal muscle cells. *Proc. Natl Acad. Sci. USA* **117**, 2978–2986 (2020).

219. Cutler, A. A., Jackson, J. B., Corbett, A. H. & Pavlath, G. K. Non-equivalence of nuclear import among nuclei in multinucleated skeletal muscle cells. *J. Cell Sci.* **131**, jcs207670 (2018).
220. Masschelein, E. et al. Exercise promotes satellite cell contribution to myofibers in a load-dependent manner. *Skelet. Muscle* **10**, 21 (2020).
221. Borowik, A. K. et al. Skeletal muscle nuclei in mice are not post-mitotic. *Function* **4**, zqac059 (2023).
This study provides the first compelling evidence that myonuclei can synthesize DNA and that this process can be augmented by muscle mechanical overload.
222. Pearen, M. A. et al. The nuclear receptor, Nor-1, markedly increases type II oxidative muscle fibers and resistance to fatigue. *Mol. Endocrinol.* **26**, 372–384 (2012).
223. Correia, J. C., Ferreira, D. M. & Ruas, J. L. Intercellular: local and systemic actions of skeletal muscle PGC-1s. *Trends Endocrinol. Metab.* **26**, 305–314 (2015).
224. Goode, J. M. et al. The nuclear receptor, Nor-1, induces the physiological responses associated with exercise. *Mol. Endocrinol.* **30**, 660–676 (2016).
225. Wu, Z. et al. Transducer of regulated CREB-binding proteins (TORCs) induce PGC-1 α transcription and mitochondrial biogenesis in muscle cells. *Proc. Natl Acad. Sci. USA* **103**, 14379–14384 (2006).
226. Wilson, T. E., Fahrner, T. J., Johnstone, M. & Milbrandt, J. Identification of the DNA binding site for NGFI-B by genetic selection in yeast. *Science* **252**, 1296–1300 (1991).
227. Maira, M., Martens, C., Philips, A. & Drouin, J. Heterodimerization between members of the Nur subfamily of orphan nuclear receptors as a novel mechanism for gene activation. *Mol. Cell Biol.* **19**, 7549–7557 (1999).
228. Chao, L. C. et al. Skeletal muscle Nur77 expression enhances oxidative metabolism and substrate utilization. *J. Lipid Res.* **53**, 2610–2619 (2012).
229. Ruas, J. L. et al. A PGC-1 α isoform induced by resistance training regulates skeletal muscle hypertrophy. *Cell* **151**, 1319–1331 (2012).
This is the first report that PGC1A4 is transcribed from the alternative PGC1A promoter after resistance exercise and can stimulate muscle hypertrophy.
230. Barres, R. et al. Acute exercise remodels promoter methylation in human skeletal muscle. *Cell Metab.* **15**, 405–411 (2012).
231. Perez-Schindler, J. et al. RNA-bound PGC-1 α controls gene expression in liquid-like nuclear condensates. *Proc. Natl Acad. Sci. USA* **118**, e2105951118 (2021).
232. Fan, W. et al. PPAR δ promotes running endurance by preserving glucose. *Cell Metab.* **25**, 1186–1193.e4 (2017).
233. Fan, L. et al. Transcription factors KLF15 and PPAR δ cooperatively orchestrate genome-wide regulation of lipid metabolism in skeletal muscle. *J. Biol. Chem.* **298**, 101926 (2022).
234. Mammucari, C. et al. The mitochondrial calcium uniporter controls skeletal muscle trophism in vivo. *Cell Rep.* **10**, 1269–1279 (2015).
235. Koh, J. H. et al. Enhancement of anaerobic glycolysis — a role of PGC-1 α 4 in resistance exercise. *Nat. Commun.* **13**, 2324 (2022).
236. Wackerhage, H. et al. Does a hypertrophying muscle fibre reprogramme its metabolism similar to a cancer cell? *Sports Med.* **52**, 2569–2578 (2022).
237. Bohlen, J., Roiuk, M. & Teleman, A. A. Phosphorylation of ribosomal protein S6 differentially affects mRNA translation based on ORF length. *Nucleic Acids Res.* **49**, 13062–13074 (2021).
238. Chaillou, T., Zhang, X. & McCarthy, J. J. Expression of muscle-specific ribosomal protein L3-like impairs myotube growth. *J. Cell. Physiol.* **231**, 1894–1902 (2016).
239. Granata, C., Oliveira, R. S. F., Little, J. P. & Bishop, D. J. Forty high-intensity interval training sessions blunt exercise-induced changes in the nuclear protein content of PGC-1 α and p53 in human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* **318**, E224–E236 (2020).
240. Brook, M. S. et al. Synchronous deficits in cumulative muscle protein synthesis and ribosomal biogenesis underlie age-related anabolic resistance to exercise in humans. *J. Physiol.* **594**, 7399–7417 (2016).
241. Norrbom, J. M. et al. A HIF-1 signature dominates the attenuation in the human skeletal muscle transcriptional response to high-intensity interval training. *J. Appl. Physiol.* **132**, 1448–1459 (2022).
242. Stokes, T. et al. Molecular transducers of human skeletal muscle remodeling under different loading states. *Cell Rep.* **32**, 107980 (2020).
243. Porter, C., Reidy, P. T., Bhattarai, N., Sidossis, L. S. & Rasmussen, B. B. Resistance exercise training alters mitochondrial function in human skeletal muscle. *Med. Sci. Sports Exerc.* **47**, 1922–1931 (2015).
244. Granata, C., Jammnick, N. A. & Bishop, D. J. Training-induced changes in mitochondrial content and respiratory function in human skeletal muscle. *Sports Med.* **48**, 1809–1828 (2018).
245. Andrade-Souza, V. A. et al. Exercise twice-a-day potentiates markers of mitochondrial biogenesis in men. *FASEB J.* **34**, 1602–1619 (2020).
246. Soto, I. et al. Balanced mitochondrial and cytosolic translatomes underlie the biogenesis of human respiratory complexes. *Genome Biol.* **23**, 170 (2022).
247. Liang, X. et al. Exercise inducible lactate dehydrogenase B regulates mitochondrial function in skeletal muscle. *J. Biol. Chem.* **291**, 25306–25318 (2016).
248. Thomas, A. C. Q. et al. Short-term aerobic conditioning prior to resistance training augments muscle hypertrophy and satellite cell content in healthy young men and women. *FASEB J.* **36**, e22500 (2022).
249. Binet, E. R. et al. Sex-based comparisons of muscle cellular adaptations after 10 weeks of progressive resistance training in middle-aged adults. *J. Appl. Physiol.* **134**, 116–129 (2023).
250. Verdijk, L. B., Snijders, T., Holloway, T. M., Van Kranenburg, J. & Van Loon, L. J. C. Resistance training increases skeletal muscle capillarization in healthy older men. *Med. Sci. Sports Exerc.* **48**, 2157–2164 (2016).
251. Hetlelid, K. J., Plews, D. J., Herold, E., Laursen, P. B. & Seiler, S. Rethinking the role of fat oxidation: substrate utilisation during high-intensity interval training in well-trained and recreationally trained runners. *BMJ Open Sport Exerc. Med.* **1**, e000047 (2015).
252. Gehlert, S. et al. Effects of acute and chronic resistance exercise on the skeletal muscle metabolome. *Metabolites* **12**, 445 (2022).
253. Ozaki, H., Loenneke, J. P., Thiebaud, R. S. & Abe, T. Resistance training induced increase in VO $_2$ max in young and older subjects. *Eur. Rev. Aging Phys. Act.* **10**, 107–116 (2013).
254. Eihara, Y. et al. Heavy resistance training versus plyometric training for improving running economy and running time trial performance: a systematic review and meta-analysis. *Sports Med. Open* **8**, 138 (2022).
255. Damas, F. et al. Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. *J. Physiol.* **594**, 5209–5222 (2016).
256. Saleem, A. & Hood, D. A. Acute exercise induces tumour suppressor protein p53 translocation to the mitochondria and promotes a p53-Tfam-mitochondrial DNA complex in skeletal muscle. *J. Physiol.* **591**, 3625–3636 (2013).
257. Beyfuss, K., Erlich, A. T., Triolo, M. & Hood, D. A. The role of p53 in determining mitochondrial adaptations to endurance training in skeletal muscle. *Sci. Rep.* **8**, 14710 (2018).
258. Li, X. et al. Ultrasensitive sensors reveal the spatiotemporal landscape of lactate metabolism in physiology and disease. *Cell Metab.* **35**, 200–211.e9 (2023).
259. Martin, A. A. et al. Sarcomere dynamics revealed by a myofibril-integrated FRET-based biosensor in live skeletal muscle fibers. *Sci. Rep.* **12**, 18116 (2022).
260. Fentz, J. et al. AMPK α is essential for acute exercise-induced gene responses but not for exercise training-induced adaptations in mouse skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* **309**, E900–E914 (2015).
261. Ballmann, C., Tang, Y., Bush, Z. & Rowe, G. C. Adult expression of PGC-1 α and - β in skeletal muscle is not required for endurance exercise-induced enhancement of exercise capacity. *Am. J. Physiol. Endocrinol. Metab.* **311**, E928–E938 (2016).
262. Lu, T., Ang, C. E. & Zhuang, X. Spatially resolved epigenomic profiling of single cells in complex tissues. *Cell* **185**, 4448–4464.e17 (2022).
263. Gandin, V. et al. Cap-dependent translation initiation monitored in living cells. *Nat. Commun.* **13**, 6558 (2022).
264. Zhen, K. et al. A systematic review and meta-analysis on effects of aerobic exercise in people with Parkinson's disease. *NPJ Parkinsons Dis.* **8**, 146 (2022).
265. Yu, Q. et al. Comparative effectiveness of multiple exercise interventions in the treatment of mental health disorders: a systematic review and network meta-analysis. *Sports Med. Open* **8**, 135 (2022).
266. Battista, F. et al. Effect of exercise on cardiometabolic health of adults with overweight or obesity: focus on blood pressure, insulin resistance, and intrahepatic fat — a systematic review and meta-analysis. *Obes. Rev.* **22**, e13269 (2021).
267. Nielsen, J. et al. Subcellular localization-dependent decrements in skeletal muscle glycogen and mitochondria content following short-term disuse in young and old men. *Am. J. Physiol. Endocrinol. Metab.* **299**, E1053–E1060 (2010).
268. Nielsen, J. et al. Increased subsarcolemmal lipids in type 2 diabetes: effect of training on localization of lipids, mitochondria, and glycogen in sedentary human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* **298**, E706–E713 (2010).
269. Yi, J. et al. Mitochondrial calcium uptake regulates rapid calcium transients in skeletal muscle during excitation-contraction (E-C) coupling. *J. Biol. Chem.* **286**, 32436–32443 (2011).
270. De Stefani, D., Raffaello, A., Teardo, E., Szabo, I. & Rizzuto, R. A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature* **476**, 336–340 (2011).
271. Baughman, J. M. et al. Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter. *Nature* **476**, 341–345 (2011).
(This reference and De Stefani et al. (2011) co-identify the core component of the elusive mitochondrial calcium uniporter.)
272. Denton, R. M., McCormack, J. G. & Edgell, N. J. Role of calcium ions in the regulation of intramitochondrial metabolism. Effects of Na $^+$, Mg $^{2+}$ and ruthenium red on the Ca $^{2+}$ -stimulated oxidation of oxoglutarate and on pyruvate dehydrogenase activity in intact rat heart mitochondria. *Biochem. J.* **190**, 107–117 (1980).
273. Turkan, A., Hiromasa, Y. & Roche, T. E. Formation of a complex of the catalytic subunit of pyruvate dehydrogenase phosphatase isoform 1 (PDP1c) and the L2 domain forms a Ca $^{2+}$ binding site and captures PDP1c as a monomer. *Biochemistry* **43**, 15073–15085 (2004).
274. Rutter, G. A. & Denton, R. M. The binding of Ca $^{2+}$ ions to pig heart NAD $^+$ -isocitrate dehydrogenase and the 2-oxoglutarate dehydrogenase complex. *Biochem. J.* **263**, 453–462 (1989).
275. Denton, R. M., Pullen, T. J., Armstrong, C. T., Heesom, K. J. & Rutter, G. A. Calcium-insensitive splice variants of mammalian E1 subunit of 2-oxoglutarate dehydrogenase complex with tissue-specific patterns of expression. *Biochem. J.* **473**, 1165–1178 (2016).
276. Sembrowich, W. L., Quintinskie, J. J. & Li, G. Calcium uptake in mitochondria from different skeletal muscle types. *J. Appl. Physiol.* **59**, 137–141 (1985).
277. Sidossis, L. S., Gastaldelli, A., Klein, S. & Wolfe, R. R. Regulation of plasma fatty acid oxidation during low- and high-intensity exercise. *Am. J. Physiol.* **272**, E1065–E1070 (1997).
278. Wang, Y. et al. Saturation of the mitochondrial NADH shuttles drives aerobic glycolysis in proliferating cells. *Mol. Cell* **82**, 3270–3283.e9 (2022).
279. Jäger, S., Handschin, C., St-Pierre, J. & Spiegelman, B. M. AMP-activated protein kinase (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1 α . *Proc. Natl Acad. Sci. USA* **104**, 12017–12022 (2007).

280. Arany, Z. et al. HIF-independent regulation of VEGF and angiogenesis by the transcriptional coactivator PGC-1 α . *Nature* **451**, 1008–1012 (2008).
281. Miyake, T. & McDermott, J. C. Re-organization of nucleolar architecture in myogenic differentiation. *J. Cell Sci.* **136**, jcs260496 (2023).
282. Emilio, E. J., Hita-Conterras, F., Jimenez-Lara, P. M., Latorre-Roman, P. & Martinez-Amat, A. The association of flexibility, balance, and lumbar strength with balance ability: risk of falls in older adults. *J. Sports Sci. Med.* **13**, 349–357 (2014).
283. Roberts, B. M., Nuckols, G. & Krieger, J. W. Sex differences in resistance training: a systematic review and meta-analysis. *J. Strength. Cond. Res.* **34**, 1448–1460 (2020).
284. Schoenfeld, B. J., Ogborn, D. & Krieger, J. W. Dose-response relationship between weekly resistance training volume and increases in muscle mass: a systematic review and meta-analysis. *J. Sports Sci.* **35**, 1073–1082 (2017).
285. Baz-Valle, E., Balsalobre-Fernandez, C., Alix-Fages, C. & Santos-Concejero, J. A systematic review of the effects of different resistance training volumes on muscle hypertrophy. *J. Hum. Kinet.* **81**, 199–210 (2022).
286. Schoenfeld, B. J., Grgic, J. & Krieger, J. How many times per week should a muscle be trained to maximize muscle hypertrophy? A systematic review and meta-analysis of studies examining the effects of resistance training frequency. *J. Sports Sci.* **37**, 1286–1295 (2019).
287. Carvalho, L. et al. Muscle hypertrophy and strength gains after resistance training with different volume-matched loads: a systematic review and meta-analysis. *Appl. Physiol. Nutr. Metab.* **47**, 357–368 (2022).
288. Refalo, M. C., Helms, E. R., Trexler, E. T., Hamilton, D. L. & Fyfe, J. J. Influence of resistance training proximity-to-failure on skeletal muscle hypertrophy: a systematic review with meta-analysis. *Sports Med.* **53**, 649–665 (2023).
289. Plotkin, D. et al. Progressive overload without progressing load? The effects of load or repetition progression on muscular adaptations. *PeerJ* **10**, e14142 (2022).
290. Bohm, S., Mersmann, F. & Arampatzis, A. Human tendon adaptation in response to mechanical loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults. *Sports Med. Open* **1**, 7 (2015).
291. Nunes, E. A. et al. Systematic review and meta-analysis of protein intake to support muscle mass and function in healthy adults. *J. Cachexia Sarcopenia Muscle* **13**, 795–810 (2022).
292. Mattioni Maturana, F., Martus, P., Zipfel, S. & AM, N. I. Effectiveness of HIIE versus MICT in improving cardiometabolic risk factors in health and disease: a meta-analysis. *Med. Sci. Sports Exerc.* **53**, 559–573 (2021).
293. Milanovic, Z., Sporis, G. & Weston, M. Effectiveness of high-intensity interval training (HIT) and continuous endurance training for VO_{2max} improvements: a systematic review and meta-analysis of controlled trials. *Sports Med.* **45**, 1469–1481 (2015).
294. Scribbans, T. D., Vecsey, S., Hankinson, P. B., Foster, W. S. & Gurd, B. J. The effect of training intensity on VO_{2max} in young healthy adults: a meta-regression and meta-analysis. *Int. J. Exerc. Sci.* **9**, 230–247 (2016).
295. Jamnick, N. A., Pettitt, R. W., Granata, C., Pyne, D. B. & Bishop, D. J. An examination and critique of current methods to determine exercise intensity. *Sports Med.* **50**, 1729–1756 (2020).
296. Iannetta, D. et al. A critical evaluation of current methods for exercise prescription in women and men. *Med. Sci. Sports Exerc.* **52**, 466–473 (2020).
297. Hov, H. et al. Aerobic high-intensity intervals are superior to improve $\dot{V}O_{2max}$ compared with sprint intervals in well-trained men. *Scand. J. Med. Sci. Sports* **33**, 146–159 (2023).
298. Hubal, M. J. et al. Variability in muscle size and strength gain after unilateral resistance training. *Med. Sci. Sports Exerc.* **37**, 964–972 (2005).
- Similar to Bouchard et al. (1999), this study clearly illustrates interindividual heterogeneity in exercise adaptation to a given resistance training programme.**
299. Volllaard, N. B. et al. Systematic analysis of adaptations in aerobic capacity and submaximal energy metabolism provides a unique insight into determinants of human aerobic performance. *J. Appl. Physiol.* **106**, 1479–1486 (2009).
300. Bouchard, C. et al. Familial aggregation of VO_{2max} response to exercise training: results from the HERITAGE Family Study. *J. Appl. Physiol.* **87**, 1003–1008 (1999).
- This is a seminal work on the contribution of genetics towards endurance training-induced changes in aerobic fitness.**
301. Montero, D. & Lundby, C. Refuting the myth of non-response to exercise training: ‘non-responders’ do respond to higher dose of training. *J. Physiol.* **595**, 3377–3387 (2017).
302. Ross, R., de Lannoy, L. & Stotz, P. J. Separate effects of intensity and amount of exercise on interindividual cardiorespiratory fitness response. *Mayo Clin. Proc.* **90**, 1506–1514 (2015).
303. Janssen, I., Heymsfield, S. B., Wang, Z. M. & Ross, R. Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J. Appl. Physiol.* **89**, 81–88 (2000).
304. Tian, Q. et al. Muscle mitochondrial energetics predicts mobility decline in well-functioning older adults: the baltimore longitudinal study of aging. *Aging Cell* **21**, e13552 (2022).
305. Alcazar, J., Rodriguez-Lopez, C., Delecluse, C., Thomis, M. & Van Roie, E. Ten-year longitudinal changes in muscle power, force, and velocity in young, middle-aged, and older adults. *J. Cachexia Sarcopenia Muscle* **14**, 1019–1032 (2023).
306. Grosicki, G. J., Zepeda, C. S. & Sundberg, C. W. Single muscle fibre contractile function with ageing. *J. Physiol.* **600**, 5005–5026 (2022).
307. Zhang, X. et al. Characterization of cellular senescence in aging skeletal muscle. *Nat. Aging* **2**, 601–615 (2022).
308. Moiseeva, V. et al. Senescence atlas reveals an aged-like inflamed niche that blunts muscle regeneration. *Nature* **613**, 169–178 (2023).
309. Dungan, C. M. et al. Deletion of SA β -Gal⁺ cells using senolytics improves muscle regeneration in old mice. *Aging Cell* **21**, e13528 (2022).
310. Dungan, C. M. et al. Senolytic treatment rescues blunted muscle hypertrophy in old mice. *Geroscience* **44**, 1925–1940 (2022).
311. Englund, D. A. et al. p21 induces a senescence program and skeletal muscle dysfunction. *Mol. Metab.* **67**, 101652 (2023).
312. Sailani, M. R. et al. Lifelong physical activity is associated with promoter hypomethylation of genes involved in metabolism, myogenesis, contractile properties and oxidative stress resistance in aged human skeletal muscle. *Sci. Rep.* **9**, 3272 (2019).
313. Ringholm, S. et al. Impact of aging and lifelong exercise training on mitochondrial function and network connectivity in human skeletal muscle. *J. Gerontol. A Biol. Sci. Med. Sci.* **78**, 373–383 (2023).
314. Rupple, B. A. et al. Resistance training rejuvenates the mitochondrial methylome in aged human skeletal muscle. *FASEB J.* **35**, e21864 (2021).
315. Vinel, C. et al. The exerkin apelin reverses age-associated sarcopenia. *Nat. Med.* **24**, 1360–1371 (2018).
316. Lee, U. et al. A Tead1-Apelin axis directs paracrine communication from myogenic to endothelial cells in skeletal muscle. *iScience* **25**, 104589 (2022).
317. O’Bryan, S. J. et al. Progressive resistance training for concomitant increases in muscle strength and bone mineral density in older adults: a systematic review and meta-analysis. *Sports Med.* **52**, 1939–1960 (2022).
318. D’Hulst, G. et al. PHD1 controls muscle mTORC1 in a hydroxylation-independent manner by stabilizing leucyl tRNA synthetase. *Nat. Commun.* **11**, 174 (2020).
319. Balachandran, A. T. et al. Comparison of power training vs traditional strength training on physical function in older adults: a systematic review and meta-analysis. *JAMA Netw. Open* **5**, e2211623 (2022).
320. McKenna, M. J., Morton, J., Selig, S. E. & Snow, R. J. Creatine supplementation increases muscle total creatine but not maximal intermittent exercise performance. *J. Appl. Physiol.* **87**, 2244–2252 (1999).
321. Hornemann, T. et al. Muscle-type creatine kinase interacts with central domains of the M-band proteins myomesin and M-protein. *J. Mol. Biol.* **332**, 877–887 (2003).
322. Kraft, T., Hornemann, T., Stolz, M., Nier, V. & Wallimann, T. Coupling of creatine kinase to glycolytic enzymes at the sarcomeric I-band of skeletal muscle: a biochemical study in situ. *J. Muscle Res. Cell Motil.* **21**, 691–703 (2000).
323. Perry, C. G. et al. Mitochondrial creatine kinase activity and phosphate shuttling are acutely regulated by exercise in human skeletal muscle. *J. Physiol.* **590**, 5475–5486 (2012).
324. Hultman, E., Soderlund, K., Timmons, J. A., Cederblad, G. & Greenhaff, P. L. Muscle creatine loading in men. *J. Appl. Physiol.* **81**, 232–237 (1996).
325. de Souza, E. S. A. et al. Effects of creatine supplementation on renal function: a systematic review and meta-analysis. *J. Ren. Nutr.* **29**, 480–489 (2019).
326. Lanher, C. et al. Creatine supplementation and lower limb strength performance: a systematic review and meta-analysis. *Sports Med.* **45**, 1285–1294 (2015).
327. Prokopidis, K. et al. Effects of creatine supplementation on memory in healthy individuals: a systematic review and meta-analysis of randomized controlled trials. *Nutr. Rev.* **81**, 416–427 (2023).
328. Areta, J. L. & Hopkins, W. G. Skeletal muscle glycogen content at rest and during endurance exercise in humans: a meta-analysis. *Sports Med.* **48**, 2091–2102 (2018).
329. Nielsen, J., Dubillot, P., Stausholm, M. H. & Ortenblad, N. Specific ATPases drive compartmentalized glycogen utilization in rat skeletal muscle. *J. Gen. Physiol.* **154**, e202113071 (2022).
- This study finds that specific depots of glycogen fuel different ATPases to varying extents in skeletal muscle.**
330. Vigh-Larsen, J. F. et al. The role of muscle glycogen content and localization in high-intensity exercise performance: a placebo-controlled trial. *Med. Sci. Sports Exerc.* **54**, 2073–2086 (2022).
331. Gejl, K. D. et al. Muscle glycogen content modifies SR Ca²⁺ release rate in elite endurance athletes. *Med. Sci. Sports Exerc.* **46**, 496–505 (2014).
332. Impey, S. G. et al. Fuel for the work required: a theoretical framework for carbohydrate periodization and the glycogen threshold hypothesis. *Sports Med.* **48**, 1031–1048 (2018).
333. Marquet, L. A. et al. Enhanced endurance performance by periodization of carbohydrate intake: “sleep low” strategy. *Med. Sci. Sports Exerc.* **48**, 663–672 (2016).
334. Tarnopolsky, M. A. et al. Influence of endurance exercise training and sex on intramyocellular lipid and mitochondrial ultrastructure, substrate use, and mitochondrial enzyme activity. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **292**, R1271–R1278 (2007).
335. Goodpaster, B. H., He, J., Watkins, S. & Kelley, D. E. Skeletal muscle lipid content and insulin resistance: evidence for a paradox in endurance-trained athletes. *J. Clin. Endocrinol. Metab.* **86**, 5755–5761 (2001).
336. Steensberg, A. et al. Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *J. Physiol.* **529**, 237–242 (2000).
337. Trinh, B. et al. Blocking endogenous IL-6 impairs mobilization of free fatty acids during rest and exercise in lean and obese men. *Cell Rep. Med.* **2**, 100396 (2021).
338. Kistner, T. M., Pedersen, B. K. & Lieberman, D. E. Interleukin 6 as an energy allocator in muscle tissue. *Nat. Metab.* **4**, 170–179 (2022).
339. Katashima, C. K. et al. Evidence for a neuromuscular circuit involving hypothalamic interleukin-6 in the control of skeletal muscle metabolism. *Sci. Adv.* **8**, eabm7355 (2022).

340. Takahashi, H. et al. TGF- β 2 is an exercise-induced adipokine that regulates glucose and fatty acid metabolism. *Nat. Metab.* **1**, 291–303 (2019).
341. Li, V. L. et al. An exercise-inducible metabolite that suppresses feeding and obesity. *Nature* **606**, 785–790 (2022).
342. Hu, M., Nie, J., Lei, O. K., Shi, Q. & Kong, Z. Acute effect of high-intensity interval training versus moderate-intensity continuous training on appetite perception: a systematic review and meta-analysis. *Appetite* **182**, 106427 (2023).
343. Reddy, A. et al. pH-gated succinate secretion regulates muscle remodeling in response to exercise. *Cell* **183**, 62–75.e17 (2020).
344. Agudelo, L. Z. et al. Skeletal muscle PGC-1 α 1 modulates kynurenine metabolism and mediates resilience to stress-induced depression. *Cell* **159**, 33–45 (2014).
345. Agudelo, L. Z. et al. Kynurenic acid and Gpr35 regulate adipose tissue energy homeostasis and inflammation. *Cell Metab.* **27**, 378–392.e5 (2018).
346. Correia, J. C. et al. Muscle-secreted neurturin couples myofiber oxidative metabolism and slow motor neuron identity. *Cell Metab.* **33**, 2215–2230.e8 (2021).

Acknowledgements

The authors apologize to all colleagues whose work could not be included owing to space constraints. The authors thank M. Karlén for preparation of the artwork in the supplementary figures. K.A.M. was supported by the National Institutes of Health (NIH R00 AG063994). K.A.D. was supported by Deutsches Zentrum für Diabetesforschung (2020/21). J.R.Z. was supported by the Swedish Research Council (Vetenskapsrådet) (2015-00165), the Swedish Research Council for Sport Science (P2022-0013, P2023-0093) and the Novo Nordisk Foundation Center for Basic Metabolic Research at the University of Copenhagen (NNF18CC0034900).

Author contributions

The authors contributed equally to all aspects of the article.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41580-023-00606-x>.

Peer review information *Nature Reviews Molecular Cell Biology* thanks Kamala Sreekumaran Nair, who co-reviewed with Mark Pataky, Adam P Sharples and Brian Glancy for their contribution to the peer review of this work.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2023, corrected publication 2023